

# DYANAP®

KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

## First Aid Treatment

**SYMPTOMS OF POISONING:** Excessive Fatigue, Sweating, Thirst and Fever. If symptoms of poisoning develop from any type of exposure to DYANAP SEND FOR A PHYSICIAN.

**FIRST AID:** Have patient lie quiet in coolest spot available. If feverish, cool with cold compresses or by immersion in cool water.

**ANTIDOTE:** If swallowed SEND FOR A PHYSICIAN. Induce vomiting by giving an emetic such as 2 tablespoonfuls of table salt in a glass of warm water, repeat until vomit fluid is clear, then give two teaspoonfuls of baking soda in a glass of water. Treat as in FIRST AID, above.


**If Splashed in Eyes,** immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.

**If Spilled on Skin,** immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If symptoms of poisoning develop, send for a physician and treat as in FIRST AID, above. Discard contaminated clothing and shoes, or clean them thoroughly before re-use. Staining may result if contaminated articles are allowed to contact carpets, upholstery, etc.

**NOTE TO ATTENDING PHYSICIAN:** Active ingredient is a metabolic stimulant. Treat symptomatically.

**IMPORTANT NOTICE:** Do not use roots, that are not recommended for this use. Do not use in areas where there are known water courses, streams, ponds, or other bodies of water. Do not use in areas where there are known water courses, streams, ponds, or other bodies of water. Do not use in areas where there are known water courses, streams, ponds, or other bodies of water.

Lot No.



KEEP OUT OF REACH OF CHILDREN  
DO NOT GET IN EYES, ON SKIN, ON CLOTHING. AVOID BREATHING SPRAY DRIFT. DO NOT TAKE INTERNALLY. DO NOT WEAR CONTAMINATED CLOTHING OR SHOES. KEEP AWAY FROM HEAT AND OPEN FLAME. (See Antidote Statement on side panel)

ABSORBED THROUGH THE SKIN -  
MAY BE FATAL IF SWALLOWED -

# DYANAP®


KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

(ALANAP®-DINITRO)  
ONE LIQUID GALLON  
PREEMERGENCE WEED KILLER

**COMPOSITION**

Active Ingredients % by weight:	
Sodium N-1-naphthyl-phthalamate*	22.3%
Sodium 4, 6-Dinitro-o-sec butyphenate**	11.5%
Inert Ingredients	66.2%
Total	100.0%

\* Equivalent to 20.8% N-1-naphthyl-phthalamic acid.  
\*\* Equivalent to 10.5% 4, 6-Dinitro-o-sec butyphenol.  
Each gallon contains two pounds of active Alanap and one pound of active DNBP.



another product of  
**UNIROYAL**

# DYANAP®

KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

## GENERAL INSTRUCTIONS

- 1. Preparation of Soil.**  
The soil should be prepared in the usual manner for the crop to be planted. The soil should be worked to a depth of 4 to 6 inches and then the DYANAP should be applied. The soil should be worked to a depth of 4 to 6 inches and then the DYANAP should be applied.
  - 2. Method of Application.**  
The DYANAP should be applied in the usual manner for the crop to be planted. The soil should be worked to a depth of 4 to 6 inches and then the DYANAP should be applied.
- Container Storage and Handling:** The container should be stored in a cool, dry place and should be kept away from heat and open flame. The container should be kept away from heat and open flame.
- Container Disposal:** The container should be disposed of in the usual manner for the crop to be planted. The soil should be worked to a depth of 4 to 6 inches and then the DYANAP should be applied.
- Equipment Cleaning:** The equipment should be cleaned in the usual manner for the crop to be planted. The soil should be worked to a depth of 4 to 6 inches and then the DYANAP should be applied.
- WARNING:** Do not use in areas where there are known water courses, streams, ponds, or other bodies of water. Do not use in areas where there are known water courses, streams, ponds, or other bodies of water.

**CAUTION: STORE ABOVE 32°F  
SHAKE WELL BEFORE USING!**

KEEP OUT OF REACH OF CHILDREN

AVOID CONTACT WITH SKIN, ON CLOTHING AVOID CONTACT WITH EYES. DO NOT TAKE INTERNALLY. WASH EXPOSED CLOTHING OR SHOES BEFORE WEARING AND OPEN FLAME. (See side panel)

# DYANAP

KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

# DYANAP

KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

# DYANAP

KILLS ANNUAL WEEDS AND GRASSES IN PEANUTS AND SOYBEANS

### GENERAL INSTRUCTIONS

#### 1. Preparation of Soil.

Soil should be prepared as usual for the crop to be planted. The soil should be free of weeds and grasses. The soil should be prepared as usual for the crop to be planted. The soil should be free of weeds and grasses.

#### 2. Method of Application.

Use DYANAP as an overall spray (broadcast) or band application. Treatment may be used over the entire area or in bands as desired. The soil should be prepared as usual for the crop to be planted.

**Container Storage and Handling.** DYANAP should be stored in a cool, dry place. The container should be kept tightly closed. The container should be kept tightly closed.

**Container Disposal.** The container should be disposed of in a safe manner. The container should be disposed of in a safe manner.

**Equipment Cleaning.** The equipment should be cleaned thoroughly after use. The equipment should be cleaned thoroughly after use.

**WARNING.** DYANAP is a potent herbicide. It should be used with care. DYANAP is a potent herbicide. It should be used with care.

**CAUTION: STORE ABOVE 32°F!  
SHAKE WELL BEFORE USING!**

**Annual Grasses.** Annual grasses are killed by DYANAP. Annual grasses are killed by DYANAP.

**Broadleaf Weeds.** Broadleaf weeds are killed by DYANAP. Broadleaf weeds are killed by DYANAP.

**DIRECTIONS FOR USE** — Put one-half the recommended amount of water in the spray tank, add the required amount of DYANAP, then add the remainder of the water to the tank while mixing. Stir or use by-pass agitation until completely mixed. It is essential that the sprayer be calibrated accurately prior to use on crop.

#### PEANUTS

Use DYANAP as an overall spray (broadcast) or band application. Treatment may be used over the entire area or in bands as desired. The soil should be prepared as usual for the crop to be planted.

**Recommended Dosage Rate for Peanuts**—Use DYANAP at the rate of 6 quarts in 20 to 40 gallons of water per acre as an overall spray (broadcast application). Treatment may be used over the entire area or in bands as desired.

Amount of DYANAP per acre to use for band treatment of Peanuts

WIDTH OF BAND	ROW SPACING					
	24 Inches	28 Inches	30 Inches	32 Inches	34 Inches	36 Inches
12"	6 Pints	5 Pints	4 1/2 Pints	4 1/2 Pints	4 1/2 Pints	4 Pints
14"	7 Pints	6 Pints	5 1/2 Pints	5 1/2 Pints	5 Pints	4 1/2 Pints

Use in 12-20 gallons of water per acre

- Note 1. Plant in a furrow slightly raised.
- Note 2. Soil should be moist at the time of application.
- Note 3. Must be used in the recommended amount.

#### SOYBEANS

Use DYANAP as an overall spray (broadcast) or band application. Treatment may be used over the entire area or in bands as desired. The soil should be prepared as usual for the crop to be planted.

**Recommended Dosage Rate for Soybeans**—Use DYANAP at the rate of 6 quarts in 20 to 40 gallons of water per acre as an overall spray (broadcast application). Treatment may be used over the entire area or in bands as desired.

Amount of DYANAP per acre to use for band treatment of Soybeans

WIDTH OF BAND	ROW SPACING			
	28 Inches	30 Inches	36 Inches	40 Inches
12"	5 Pints	4 1/2 Pints	4 Pints	3 1/2 Pints
14"	6 Pints	5 1/2 Pints	4 1/2 Pints	4 Pints

Use in 9 gallons of water per 40 inches with 1 1/2 quarts of water per inch of row

- Note 1. Plant in a furrow slightly raised.
- Note 2. Soil should be moist at the time of application.
- Note 3. Must be used in the recommended amount.

**DO NOT USE DYANAP ON SILT LOAM**

**ACCEPTED**  
JAN 13 1972  
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND HERBICIDE ACT FOR ECONOMIC POISON CONTROL. REGISTERED UNDER NO. 400-75

another product of  
**UNIROYAL**

UNION CARBIDE CORPORATION  
PO BOX 1170  
PHILADELPHIA, PA 19106