

DANGER! KEEP OUT OF REACH OF CHILDREN.

Do not get in eyes.

Causes severe eye damage.

Wear goggles when handling.

In case of contact with eyes, flush immediately with running water and get medical attention.

For other danger statements see below.

VITAVAX[®] FUNGICIDE

Net Weight 50 Lbs.

COMPOSITION

| | |
|----------------------------------------------------------------------------------|-----|
| Active Ingredient: Carboxin (5,6-dihydro-2-methyl-1,4-oxathin-3-carboxanilide) * | 75 |
| Inert Ingredients: | 25 |
| Total: | 100 |

* U.S. Patent Nos. 3,249,499 - 3,393,202 - 3,454,391

1972 UNIROYAL, INC. —
All rights reserved.
EPA Reg. No. 400-80-AA

E.P.A. Est. No.



DIRECTIONS FOR USE

VITAVAX Fungicide is a wettable powder seed treatment for use as a dry dust, spray-mist, or slurry seed treatment. In liquid systems, a thin paste should first be made using a portion of the required water and the recommended amount of VITAVAX Fungicide. Stir this paste into the remainder of the water and provide enough agitation to maintain a uniform water suspension.

COTTON

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases of cotton. Seed rot, damping off and sore-shin due to *Rhizoctonia solani* are prevented. VITAVAX may be used on both machine and acid delinted cottonseed. Use 8 ounces of VITAVAX Fungicide per 100 pounds of seed. On seed previously treated with the recommended rate of a standard fungicide such as captan or thiram, 4 ounces of VITAVAX Fungicide may be used as an overtreatment.

USE RESTRICTIONS — Do not use treated cottonseed for food, feed or oil purposes. Do not graze or feed livestock on hay grown from treated seed.

COMPATIBILITY — VITAVAX Fungicide can be used in combination with other seed protectant fungicides such as captan or thiram. It can be used in com-

USE RESTRICTIONS — Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

COMPATIBILITY — VITAVAX Fungicide is compatible with seed protectant fungicides such as thiram.

PEANUTS

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases on peanuts. It is particularly effective against *Rhizoctonia solani*. Use 3 to 6 ounces of VITAVAX Fungicide per 100 pounds of seed. VITAVAX Fungicide can also be used at 2 to 3 ounces per 100 pounds of seed when used with seed protectant fungicides such as captan, thiram or Difoatan.

USE RESTRICTIONS — Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas. Do not hog down treated peanut fields prior to harvest.

COMPATIBILITY — VITAVAX Fungicide is compatible with seed protectant fungicides such as captan, thiram or Difoatan.

ADDITIONAL CAUTIONS AND INSTRUCTIONS

HANDLING — Harmful if swallowed. Avoid inhaling dust or spray. Wash hands and face thoroughly with soap and

VITAVAX Fungicide. Stir the paste into the remainder of the water and provide enough agitation to maintain a uniform water suspension.

COTTON

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases of cotton. Seed rot, damping off and sore-shin due to *Rhizoctonia solani* are prevented. VITAVAX may be used on both machine and acid delinted cottonseed. Use 8 ounces of VITAVAX Fungicide per 100 pounds of seed. On seed previously treated with the recommended rate of a standard fungicide such as captan or thiram, 4 ounces of VITAVAX Fungicide may be used as an overtreatment.

USE RESTRICTIONS — Do not use treated cottonseed for food, feed or oil purposes. Do not graze or feed livestock on hay grown from treated seed.

COMPATIBILITY — VITAVAX Fungicide can be used in combination with other seed protectant fungicides such as captan or thiram. It can be used in combination with seed treatment insecticides such as Di-Syston or Thimet.

BARLEY, OATS and WHEAT

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases of barley, oats and wheat. It is particularly effective against *Rhizoctonia solani*.

In addition, loose smut, covered smut, flag smut, common bunt and barley stripe are controlled by VITAVAX Fungicide.

ACCEPTED
SEP 11 1975
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-80

DOSAGE INSTRUCTIONS

| Crop | Diseases Controlled | Amount of VITAVAX Fungicide per 100 pounds of seed |
|--------|-------------------------------------------------------------------------------|----------------------------------------------------|
| Barley | seed rots seedling diseases barley stripe loose smut covered smut | 2-3 ounces** |
| Oats | seed rots seedling diseases loose smut covered smut | 1-1.5 ounces 2 ounces |
| Wheat | seed rots seedling diseases loose smut common bunt flag smut | 2 ounces |

**NOTE: On barley and wheat seed grown for planting, seed may also be used at the rate of four ounces per 100 pounds of seed.

Lot No.

UNIROYAL CHEMICAL — DIVISION OF UNIROYAL, INC.
 Naugatuck, Connecticut 06770

PEANUTS

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases on peanuts. It is particularly effective against *Rhizoctonia solani*. Use 3 to 6 ounces of VITAVAX Fungicide per 100 pounds of seed. VITAVAX Fungicide can also be used at 2 to 3 ounces per 100 pounds of seed when used with seed protectant fungicides such as captan, thiram or Difolatan.

USE RESTRICTIONS — Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas. Do not hog down treated peanut fields prior to harvest.

COMPATIBILITY — VITAVAX Fungicide is compatible with seed protectant fungicides such as captan, thiram or Difolatan.

ADDITIONAL CAUTIONS AND INSTRUCTIONS

HANDLING — Harmful if swallowed. Avoid inhaling dust or spray. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Keep away from food or food products. Care must be exercised in the handling of treated seed.

SEED LABELING — "This seed treated with VITAVAX Fungicide. Do not use for food, feed or oil purposes." Add appropriate information for any other chemical treatments.

FISH AND WILDLIFE — Do not contaminate water by cleaning of equipment or disposal of wastes.

CONTAINER DISPOSAL — Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food or water. If disposal is by burning, keep out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.