


400-82

5-6-1971

1/5



WARNING

KEEP OUT OF REACH OF CHILDREN

(For other caution statements see back panel) . ||

OMITE®-30W
AGRICULTURAL MITICIDE

ACCEPTED
MAY 6 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR
ED WITH IN 400-82 REGISTER
AND ATTACHED SUBJECT

COMPOSITION

Active Ingredient (% by weight)	
2-(p-tert-butylphenoxy)cyclohexyl 2- propynyl sulfite*	30.0
Inert	<u>70.0</u>
Total	100.0

*U.S. Patents 3,272,854; 3,463,859

®UniRoyal's Registered Trademark for its Miticide

2/5

4-2-57
1000

OMITE-30W

A dry wettable powder for preparing sprays to control the strawberry spider, brown almond, European red, Pacific spider and two-spotted spider mites.

Omite-30W is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Mites hit by the spray stop feeding and are killed within 48 to 72 hours, depending on temperature. Omite has some ovicidal activity. The young mites that do hatch from eggs after the spray is applied will be killed by the residue of Omite left on the leaves and fruit.

Add recommended dosage of Omite-30W to the spray tank, add water, agitate and spray thoroughly to cover foliage (both upper and lower leaf surfaces) and fruit for best results. See table for dosage rates.

- Apricots (west of Rockies) For the control of two-spotted and Pacific mites.
- Nectarines (west of Rockies) For the control of Pacific and brown almond mites.
- Strawberries (west of Rockies only) For the control of Pacific Strawberry and two-spotted spider mites.
- Hops - For the control of two-spotted spider mites.

3/5
8

OMITE-30W

Compatibility - Omite-30W is compatible with captan, Chloro-benzilate, Cyprex, DDD, DDT, Diazinon*, dieldrin, Endrin, Ethion, Folpet, fixed neutral copper, glyodin, Guthion WP*, Karathane, lead arsenate, malathion, methyl parathion, parathion, Phosdrin, Phygon, Sevin, rotenone, R-meta-Systox, sabadilla, Systox, TEPP, Trithion, Thiodan, thiram, Toxaphene, zineb, zinc oxide, Dikar, ferbam, Gardona, Imidan, maneb and sulfur.

*Caution - Do not use at day temperatures above 95°F west of the Rocky Mountains, 85°F east of the Rocky Mountains.

NOTE - Do not use Omite-30W mixed with alkaline materials such as lime, lime sulfur or Bordeaux mixture. Do not use spring or summer foliar oil sprays before, with, or following Omite-30W. Do not use more than four ounces of additional spray adjuvant per 100 gallons. Do not add any additional spray adjuvants to the spray tank if Omite-30W is mixed with one or more pesticide formulations.

OMITE-30W

WARNING

Handling

Avoid inhaling dust or spray.

Do not get in eyes. May cause eye damage. The wearing of goggles is recommended. In case of contact with eyes, flush with running water for 15 minutes and get medical attention.

In case of skin contact, wash thoroughly with warm water and soap.

May be harmful if swallowed.

Fish and Wildlife

Keep out of any body of water. This product is toxic to fish.

Use Restrictions

Do not graze or feed livestock on cover crops grown among the trees and vines listed on this label.

Do not feed hop refuse or cones to livestock.

Container Disposal

Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CONTROL NO.

USDA REGISTRATION NO.

UniRoyal Chemical - Division of UniRoyal, Inc.
Naugatuck, Connecticut 06770

OMITE-30W

Crops	Mites Controlled	Amount Per Acre	Dosage Instructions		Gallons of Spray Solution Per Acre	Total Number of Sprays Per Year	Days Before Harvest (Last Spr)
			Amount per 100 Gallons of Water				
			West of Rocky Mountains	East of Rocky Mountains			
Necta- rines	Pacific spider brown almond	9 lbs.	1½ lbs.	Do not use	600	2	14
Apricots	Pacific spider two-spotted spider	6-9 lbs.	1½ lbs.	Do not use	400 - 600	2	14
Straw- berries (see note)	Pacific spider strawberry spider two-spotted spider	5-10 lbs.	5 lbs.	Do not use	100 - 200	3	3
Hops	two-spotted spider	5 lbs.	2½ lbs.	---	200	2	14

NOTE: "Apply as necessary but no more than three applications can be made per crop season."
(bloom to harvest).

5/5