

400-82

12-3-1972

1/2

**WARNING! KEEP OUT OF REACH OF CHILDREN.**

For other warning statements see back panel.



# OMITE<sup>®</sup> 30W AGRICULTURAL MITICIDE

(West of the Rocky Mountains)

ACCEPTED  
DEC 3 1972  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 400-82 SUBJECT  
TO ATTACHED COMMENTS.

NET WT. 5 POUNDS

| COMPOSITION  |      |
|--|------|
| Active ingredient (% by weight)                                |      |
| 2-(p-tert-butylphenoxy) cyclohexyl<br>2-propynyl sulfite*..... | 30%  |
| Inert ingredients.....   | 70%  |
| Total  | 100% |

\*U.S. PATENT NOS. 3,272,854 and 3,463,859  
EPA REGISTRATION NO. 400-82-AA

*another product of*



# OMITE 30W

AGRICULTURAL MITICIDE

2/2

A dry, wettable powder for preparing sprays to control European red, two-spotted spider, McDaniel, Pacific spider, peach silver, brown almond and strawberry spider mites. Omite-30W is not systemic in action; therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of Omite-30W per acre.

### DOSAGE INSTRUCTIONS

| Crop  | Mites controlled  | Amount OMITE-30W per acre per treatment | OMITE-30W per 100 gallons of water West of Rocky Mountains | Gallons spray solution per acre | Total number of sprays /year | Last spray days before harvest |
|---|---|---|--|---------------------------------|------------------------------|--------------------------------|
| APPLES  | European red<br>Two-spotted spider<br>Pacific spider<br>McDaniel                                      | 5-12 lbs.                               | 1-1 1/2 lbs.   | 500-800                         | 3                            | 7                              |
| APRICOTS                                      | Pacific spider<br>Two-spotted spider  | 6-9 lbs.                                | 1 1/2 lbs.   | 400-600                         | 2                            | 14                             |
| CALIFORNIA WALNUTS                            | European red<br>Two-spotted spider  | 7-22 lbs.                               | 1 1/2 lbs.   | 500-1500                        | 2                            | 14                             |
| HOPS  | Two-spotted spider  | 5 lbs.                                  | 2 1/2 lbs.   | 200                             | 2                            | 14                             |
| NECTARINES                                    | Pacific spider<br>Brown almond  | 9 lbs.                                  | 1 1/2 lbs.   | 600                             | 2                            | 14                             |
| PEACHES                                       | European red<br>Pacific spider<br>Two-spotted spider<br>Strawberry spider<br>McDaniel<br>Peach silver | 5-15 lbs.                               | 1 1/2 lbs.   | 300-1000                        | 2                            | 14                             |
| PLUMS<br>PRUNES                               | European red<br>Two-spotted spider<br>Pacific spider  | 5-15 lbs.                               | 1 1/2 lbs.   | 300-1000                        | 2                            | 28                             |
| STRAWBERRIES<br>(See application notes below) | Pacific spider<br>Strawberry spider<br>Two-spotted spider   | 5-10 lbs.                               | 5 lbs.   | 100-200                         | 3*                           | 3                              |

\*\*\*Apply as necessary but no more than three applications can be made per crop season.\*\*\* (bloom to harvest.)

### APPLICATION NOTES (Strawberries only)

1. Do not combine with other chemicals or spray adjuvants, as leaf and fruit injury may occur.
2. California: Do not apply at day temperatures above 80°F as fruit burn may occur.

### FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application

### DOSAGE INSTRUCTIONS

| Crop               | Mites controlled  | Amount OMITE-30W per acre per treatment | OMITE-30W per 100 gallons of water West of Rocky Mountains | Gallons spray solution per acre | Total number of sprays /year | Last spray days before harvest |
|--------------------|---|---|--|---------------------------------|------------------------------|--------------------------------|
| ALMONDS            | Clover, European red, Two-spotted spider, Strawberry spider, Pacific spider | 5-25 lbs.                               | 1 1/2 lbs.   | 500-1500                        | —                            | —                              |
| GRAPES             | Pacific spider<br>Strawberry<br>Willamette<br>Two-spotted spider            | 5-6 lbs.                                | 1 1/2 lbs.   | 300-400                         | —                            | —                              |
| ORANGES AND LEMONS | Citrus red  | 10-30 lbs.                              | FOR NON-BEARING TREES ONLY<br>1-2 lbs.                     | 1000-1500                       | —                            | —                              |
| PEARS              | European red<br>Two-spotted spider  | 5-7 1/2 lbs.                            | FOR POST HARVEST TREES ONLY<br>1 1/2 lbs.                  | 300-500                         | —                            | —                              |

APPLICATION NOTE: Omite performance is best when daily temperatures average above 70°F.

### WARNING:

**HANDLING:** Avoid inhaling dust or spray. Do not get in eyes. May cause eye damage. The wearing of goggles is recommended. In case of contact with eyes, flush with running water for 15 minutes and get medical attention. In case of skin contact, wash thoroughly with warm water and soap. May be harmful if swallowed.

**FISH AND WILDLIFE:** Keep out of any body of water. This product is toxic to fish. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

**USE RESTRICTIONS:** Do not graze or feed livestock on cover crops grown among the trees listed on this label. Do not feed animals apple pomace made from Omite treated apples. Do not feed hop refuse or cones to livestock.

### CAUTION:

A. Leaf injury and fruit spotting may occur if the following conditions are not observed:

1. Do not apply Omite-30W when air temperatures are above 95°F.
2. Do not apply Omite-30W in combination with Guthion emulsifiable concentrate.
3. Do not apply Omite-30W before, with, or following spring and summer foliar spray oils.

(Further information available from supplier.)

4. Do not use more than four ounces of additional spray adjuvant per 100 gallons if Omite-30W is used alone. Add no spray adjuvant if Omite-30W is mixed with one or more pesticide formulations in the spray tank.

B. Omite-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

**COMPATIBILITIES:** Except as noted above, Omite-30W is compatible with most pesticides. (See caution statements.) Check with your supplier for details.

**CONTAINER DISPOSAL:** Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food.

**IMPORTANT NOTICE:** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.