

400-82

5-7-1973

RECORDED  
20 JUNE 73 1/5  
JK

WARNING! KEEP OUT OF REACH OF CHILDREN  
For other warning statements see back panel

OMITE (R) -30W

Agricultural Miticide

NEW WEIGHT 5 POUNDS

ANOTHER PRODUCT OF UNIROYAL

COMPOSITION

Active Ingredient (% by weight)	
2-(p-tert-butylphenoxy)cyclohexyl 2-propynyl sulfite*	.. 30%
Inert Ingredients .....	70%
Total .....	100%

\*U.S. Patent No. 3,272,854, 3,463,859

312-67F-1169

(c) 1969 UNIROYAL, Inc. All Rights Reserved

ACCEPTED

MAY 7 - 1973

REGISTERED UNDER THE FEDERAL INSECTICIDE,  
FUNGICIDE AND RODENTICIDE ACT,  
FOR USE AS A POISON REGISTERED  
ED UNDER NO. 400-82. SUBJECT  
TO ATTACHED COMMENTS.

2/5

OMITE-30W

Agricultural Miticide

A dry wettable powder for preparing sprays to control European red, McDaniel, Pacific spider, peach silver, strawberry spider and two-spotted spider mites.

Omite-30W is not systemic in action, therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Mites hit by the spray stop feeding and are killed within 48 to 72 hours, depending on temperature. Young mites hatching from eggs after the spray is applied will be killed by the residue of Omite left on the leaves and fruit.

Add recommended dosage of Omite-30W to the spray tank, agitate and spray thoroughly to cover foliage and fruit for best results.

Almonds • California only

To control clover, European red, strawberry spider, two-spotted spider and Pacific spider mites.

Use 5 to 22 pounds Omite 30W per acre either as a dilute or concentrate spray.

As a dilute spray add 1 to 1½ pounds Omite 30W to 100 gallons of water and apply from 500 to 1,500 gallons of liquid per acre.

Two treatments can be made per season. The last spray can be applied 28 days before harvest.

OMITE-30W

Agricultural Miticide

Grapes - California only

To control strawberry spider, Willamette, Pacific spider and two-spotted spider mites.

Use 5 to 9 pounds Omite 30W per acre as a dilute spray. Add 1½ pounds Omite 30W in 100 gallons of water. Apply 350 to 600 gallons of liquid per acre.

Two treatments can be made per season. The last treatment can be made up to 21 days of harvest.

Oranges, lemons - California only

For the control of citrus red mite Panonychus citri (McG.)

Use 10 to 15 pounds Omite 30W per acre as a dilute spray. (Do not use less than 10 pounds per acre on mature trees.)

Add 1 to 1½ pounds Omite 30W to 100 gallons of water and apply 600 to 1,500 gallons of liquid per acre.

Two treatments can be made in one season, the last can be applied up to 7 days before harvest.

(Omite 30W may cause some leaf burn on new growth - (feather growth).)

WARNING

HANDLING - Avoid inhaling dust or spray. Do not get in eyes. May cause eye damage. The wearing of goggles is recommended. In case of contact with eyes, flush with running water for 15 minutes and get medical attention. In case of skin contact, wash thoroughly with warm water and soap. May be harmful if swallowed.

Omite 30W

Agricultural Miticide

FISH AND WILDLIFE - Keep out of any body of water. This product is toxic to fish.

USE RESTRICTIONS - Do not graze or feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITY - Omite-30W is compatible with captan, chlorobenzilate, C-Lite, Cygon, Cyprex, DDD, DDT, Diazinon (see note), dibrom, dieldrin, dikar, endrin, ethion, ferbam, fixed neutral copper, folpet, gardona, glyodin, guthion WP (see note), imidan, karathane, lead arsenate, malathion, maneb, methyl parathion, parathion, phosdrin, phygon, R-meta-Systox, rotenone, sabadilla, sevin,, sulfur, systox, TEPP, thiodan, thiram, toxaphene, trithion, urea, zinc oxide and zineb.

NOTES

1. Caution - Do not use in combination with diazinon or guthion WP when air temperatures are above 95°F west of the Rocky Mountains or 85°F east of the Rocky Mountains, as leaf injury may occur.
2. Do not use with guthion emulsifiable concentrate as leaf injury may occur.
3. Do not apply Omite-30W before, with or following spring and summer foliar spray oils or leaf injury and fruit spotting may occur.
4. Do not use more than four ounces of additional spray adjuvant per 100 gallons if Omite-30W is used alone or leaf injury may occur.

Omite-30W

Agricultural Miticide

Add no spray adjuvant if Omite-30W is mixed with one or more pesticide formulations in the spray tank to avoid leaf injury and fruit spotting.

5. Omite-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. Effectiveness of Omite will be reduced.

6. Omite-30W performance is best when temperature is 70°F or higher. Complete and uniform coverage is necessary for optimum results.

CONTAINER DISPOSAL - Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food.

IMPORTANT NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

LOT NO.

EPA Registration No. 400-82-AA

UNIROYAL CHEMICAL - Division of UNIROYAL, Inc. - Naugatuck, Connecticut 06770