DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage. Do not get in eyes, on skin or on clothing.

Wear goggles or face shield and rubber gloves when handling.

Harmful if swallowed. Avoid inhaling dust or spray.

Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements, see back panel.

ACCEPTED

JUL 1 5 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 100-82 EPA Reg. No.



Net Weight: 5 POUNDS

COMPOSITION

*U.S. PATENT NOS. 3,272,854 and 3,463,859 EPA REGISTRATION NO 400-82-AA

E P A Est. No. 279-CA-2



Willamette mites. Omite-30W is not systemic in action, therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of Omite-30W per acre. Omite-30W can be used in concentrate sprays.

DOSAGE INSTRUCTIONS

		Amount OMITE-30W per ocre per	OMITE-3CW per 100 gallians of water West of Rocky	Gallons spray solution	Total number al sprays	Last spray days before
Crop	Mites controlled	treatment	Mountains	per ocre	year	harvest
ALMONDS (Colifornia only)	clover, European red, two-spotted spider, strowberry spider, Pacific spider	5-22 fbs.	[-1 ² 2 (bs.:	500-1500	2	26
APPLES	European red two-spotted spider Pacific spider McDaniel	5-12 lbs.	1-1½ lbs.	500-800	3	7
APRICOTS	Pacific spider two-spotted spider	5-9 lbs.	1% 16s.	300-600	2	14
CALIFORNIA WALNUTS	European red two-spotted spider	7-22 lbs.	1½ lbs.	500-1500	2	14
FIGS	twa-spatted spider Pacific spider strawbeery spider	3≈6 lbs.	1-1% ths.	200 - 400	2	14
GRAPEFRUIT (California anly)	citrus red	15 Іъз.	1½ lbs.	1000	2	7
GRAPES (California anly)	Pacific spider strawberry spider Wilametre two-spotted spider	5-9 lbs.	1% lbs.	350-600	2	21
HOPS	two-spotted spider	5 lbs.	25i lbs.	200	2	14
NECTARINES	Pacific spider brown almond	5-9 lbs.	132 lbs.	300-600	2	14
ORANGES AND LEMONS (Coldernia only)	citrus sed	10-15 lbs.	I-1½ lbs.	600-1500	2	7
PEACHES	Europeon red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5-15 lbs.	14 160.	300-1000	2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5-15 lbs.	1½ ibs.	300-1000	2	14
ROSES (greenhouse grown)	two-spotted spider Pacific spider smawberry spider		116.			_
Crop	Mites controlled	Amount OMITE-30W per acre per treatment	OMITE-30% per 100 gallons of water West of Recky Mountains	Gallons spray sulution per acra	Total no. of sprays: crop season (bloom- harvest)	Last spray days before harvest
STRAWBERRIES	Pocific spider	5-10 lbs.	5 (bs.	100-200	3.	3

Do not apply more than 3 applications of Omite's any 40 day period during the hervesting season.

APPLICATION NOTES

(See application strawberry spider notes below) two-spotted spider

- Do not combine with other chemicals or spray adjuvants, as leef and fruit injury may occur. (strawberries only)
 <u>Colitornja</u>: Do not apply at day temperatures above 80° F as fruit burn may occur. (strawberries only)
 Omite-J0W may cause some leaf burn on new growth (feather growth) of aranges and femons.
 Omite performance is best when doily temperatures average above 70° F.

FOR POST HARVEST AND NON-BEARING TREES ONLY Nombearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

Стор	Mites controlled	Amount OMITE-30W per acre per treatment	OMITE-3CW per 100 gallans of water West of Rocky Mountairs	Gallons spray solution per ocre	Total number of sprays year	Last spray days before harvest
CHERRIES	European red two-spotted spider Pacific spider McDaniel	6 lbs.	He lbs.	400		
PEARS	European red two-spotted spider	5-7½ lbs.	FOR POST HA			

FISH AND WILDLIFE: This product is toxic to fish, shrimp and crab. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this tast?

USE RESTRICTIONS: Do not feed livestock on cover crops grown among the trees listed on this label. Do not feed hop refuse or cones to livestock. However, spent hops from the extract operation may be fed.

- A. Leaf injury and/or fruit spotting may occur if the following conditions are not observed.

 1. Do not apply Omite-30W with Diazinon-WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.

 2. Omite-30W is not compatible with foliar spray oils. (Further information available from sense).
 - from supplier.)
 3. Sprint-38* is the spreader-sticker designed for use with Omite-30W; other spreader-stickers may be used (follow suppliers' recommendation on rates).

B. Omite-30W is not compatible with alkaline materials such as time, time sulfur or Bordeaux. The effectiveness of Omite will be reduced.

COMPATIBILITIES: Except as noted above, Omite-30W is compatible with most pesticides. (See causion satements.) Check with your supplier for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CONTAINER DISPOSAL: Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is presect contours to its chemical description and is teasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions or under constitues or under con ditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

UNIRGYAL CHEMICAL - Division of UNIRGYAL, for Nauratuck Connecticut 06770



CCC