

400-82

7-30-1976

1/2

DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage.

Do not get in eyes, on skin or on clothing.

Wear goggles or face shield and rubber gloves when handling.

Harmful if swallowed. Avoid inhaling dust or spray.

Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements, see back panel.

EPA-PA

EPA-PA

OMITE[®]

30W

AGRICULTURAL MITICIDE

(West of the Rocky Mountains)

ACCEPTED**JUL 30 1976**

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 400-82

Net Weight:**COMPOSITION**

Active Ingredient (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	30%
Inert Ingredients	70%
Total	100%

*U.S. PATENT NOS. 3,272,854 and 3,463,859
EPA REGISTRATION NO. 400-82-AA

EPA Est. No. 279-CA-2**UNIROYAL**

A dry wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two-spotted spider and Wilamette mites.

Omite-30W is not systemic in action, therefore,

complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of Omite-30W per acre. Omite-30W can be used in concentrate sprays.

DOSAGE INSTRUCTIONS

Crop	Mites controlled	Amount OMITE-30W per acre per treatment	OMITE-30W per 100 gallons of water West of Rocky Mountains	Gallons spray solution per acre	Total number of sprays /year	Last spray days before harvest
ALMONDS (California only)	clover, European red, two-spotted spider, strawberry spider, Pacific spider	5-22 lbs.	1-1½ lbs.	500-1500	2	28
APPLES	European red two-spotted spider Pacific spider McDaniel	5-12 lbs.	1-1½ lbs.	500-800	3	7
APRICOTS	Pacific spider two-spotted spider	5-9 lbs.	1½ lbs.	300-600	2	14
CALIFORNIA WALNUTS	European red two-spotted spider	7-22 lbs.	1½ lbs.	500-1500	2	14
FIGS	Two-spotted spider Pacific spider strawberry spider	3-6 lbs.	1-1½ lbs.	200-400	2	14
GRAPEFRUIT (California only)	citrus red	15 lbs.	1½ lbs.	1000	2	7
GRAPES (California only)	Pacific spider strawberry spider Wilamette two-spotted spider	5-9 lbs.	1½ lbs.	350-600	2	21
HOPS	two-spotted spider	5 lbs.	2½ lbs.	200	2	14
NECTARINES	Pacific spider brown almond	5-9 lbs.	1½ lbs.	300-600	2	14
ORANGES AND LEMONS (California only)	citrus red	10-15 lbs.	1-1½ lbs.	600-1500	2	7
PEACHES	European red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5-15 lbs.	1½ lbs.	300-1000	2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5-15 lbs.	1½ lbs.	300-1000	2	14
ROSES (greenhouse grown)	two-spotted spider Pacific spider strawberry spider	—	1 lb.	—	—	—

Crop	Mites controlled	Amount OMITE-30W per acre per treatment	OMITE-30W per 100 gallons of water West of Rocky Mountains	Gallons spray solution per acre	Total no. of sprays/ crop season (bloom-harvest)	Last spray days before harvest
STRAWBERRIES (See application notes below)	Pacific spider strawberry spider two-spotted spider	5-10 lbs.	5 lbs.	100-200	3*	3

* Do not apply more than 3 applications of Omite in any 40 day period during the harvesting season.

APPLICATION NOTES

- Do not combine with other chemicals or spray adjuvants, as leaf and fruit injury may occur. (strawberries only)
- CALIFORNIA: Do not use OMITE-30W on oranges and grapefruit from the time new fruit is present in the spring until September first, as fruit spotting may occur. Do not apply at day temperatures above 80°F as fruit burn may occur (Strawberries ONLY).
- Omite-30W may cause some leaf burn on new growth (feather growth) of oranges and lemons.
- Omite performance is best when daily temperatures average above 70° F.

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

Crop	Mites controlled	Amount OMITE-30W per acre per treatment	OMITE-30W per 100 gallons of water West of Rocky Mountains	Gallons spray solution per acre	Total number of sprays /year	Last spray days before harvest
CHERRIES	European red two-spotted spider Pacific spider McDaniel	6 lbs.	1½ lbs.	400	—	—
PEARS	European red two-spotted spider	5-7½ lbs.	1½ lbs.	300-500	—	—

WARNING:

FISH AND WILDLIFE: This product is toxic to fish and shrimp. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

USE RESTRICTIONS: Do not feed livestock on cover crops grown among the trees listed on this label. Do not feed hop refuse or cones to livestock. However, spent hops from the extract operation may be fed

CAUTION:

A. Leaf injury and/or fruit spotting may occur if the following conditions are not observed.

- Do not apply Omite-30W with Diazinon-WP if air temperatures are expected to exceed 95° F the day of spraying apples and hops.
- Omite-30W is not compatible with foliar spray oils. (Further information available from supplier or Uniroyal.)
- Sprint-38® is the spreader-sticker designed for use with Omite-30W.
- Do not follow or precede an OMITE-30W application with a dimethoate application within a 21 day period, when fruit is present.

5. Do not use any additional surfactants or wetting agents with OMITE-30W.

8. Omite-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

COMPATIBILITIES: Except as noted above, Omite-30W is compatible with most pesticides. (See caution statements.) Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CONTAINER DISPOSAL: Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

UNIROYAL CHEMICAL— Division of UNIROYAL, Inc. Naugatuck, Connecticut 06770

7-21-75-W

© 1969, 1971, 1973, Uniroyal Inc. All Rights Reserved