

**PRECAUTIONARY  
STATEMENTS  
HAZARDS TO HUMANS  
DANGER**

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable, keep away from heat and open flame.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**1. PROHIBITIONS**

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

**2. PESTICIDE DISPOSAL**

Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**3. CONTAINER DISPOSAL**

Triple rinse (or equivalent) and offer for recycling, or disposal in approved landfill or bury in a safe place.

**4. GENERAL**

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

**OMITE<sup>®</sup>-30W**

AGRICULTURAL MITICIDE  
(EAST OF THE ROCKY MOUNTAINS)



**COMPOSITION**

Active Ingredient: (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	30.0%
Inert Ingredients:	70.0%
Total:	100.0%

\*U.S. Patents 3,272,854 - 3,463,859

**KEEP OUT OF REACH OF CHILDREN.  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**IF INHALED:** See a physician if an abnormal reaction occurs.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF IN EYES:** Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-82-AA

©1969, 1971 UNIROYAL, Inc. All Rights Reserved.

NET WEIGHT

10-17-77

400-82  
28-004  
11-1-1977


**DIRECTIONS FOR USE**

A dry, wettable powder for preparing sprays to control European red, two-spotted spider, McDaniel, Pacific spider, peach silver, and strawberry spider mites.

OMITE-30W is not systemic in action; therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-30W per acre.

**DOSAGE INSTRUCTIONS**

See other side for Dosage Instructions Charts.



1/2



### DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER EAST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
APPLES	European red two-spotted spider Pacific spider McDaniel	5-12 lbs.	1¼-1½ lbs.	400-800	3	7
PEACHES	European red two-spotted spider peach silver strawberry spider McDaniel Pacific spider	5-15 lbs.	1¼-1½ lbs.	400-1000	2	14
PEANUTS	two-spotted spider	3-5 lbs.	See application note 8 below		2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5-15 lbs.	1¼-1½ lbs.	400-1000	2	14
CARNATIONS CHRYSANTHEMUMS ROSES	two-spotted spider	—	1-1½ lbs.	Spray to wet	—	—

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER EAST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER CROP SEASON (BLOOM - HARVEST)	LAST SPRAY DAYS BEFORE HARVEST
STRAWBERRIES (Florida and Louisiana only)	two-spotted spider	2½	2½	100	3*	3

\*Do not apply more than 3 applications of Omite in any 21 day period during the harvesting season.

#### FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

### DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER EAST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
PEARS	European red two-spotted spider	5-7½ lbs.	1¼-1½ lbs.	400-500	—	—

1. COMPLETE AND UNIFORM COVERAGE IS NECESSARY FOR OPTIMUM RESULTS.
2. ALWAYS USE AT LEAST THE MINIMUM RECOMMENDED AMOUNT OF OMITE-30W PER ACRE.

3. Omite-30W can be used in concentrate sprays.
4. Do not reduce the rate per acre when applying concentrate sprays.
5. Omite performance is best under normal summer temperatures.
6. Do not exceed the recommended amount of OMITE-30W per acre.
7. Use on roses is limited to greenhouse, slat and saran grown. Because cultivar reaction and cultural practices differ widely, the user should initially treat only a small portion of his crop to confirm plant safety under his conditions. Repeat applications of OMITE-30W are normally necessary for extended control.
8. PEANUTS — Premix OMITE-30W with a small amount of water to form a slurry before adding to spray tank. Apply in sufficient water to provide complete coverage of foliage when using low gallonage equipment. A minimum of 20 gallons of water per acre is recommended for best results. Do not plant rotational crops within 6 months of the last treatment on peanuts.

**USE RESTRICTIONS —** Do not graze or feed livestock on cover crops grown among the trees listed on this label.

#### CAUTION:

- A. Leaf injury and fruit spotting may occur if the following conditions are not observed:
1. Do not apply OMITE-30W when air temperatures are above 90°F.
  2. Do not apply OMITE-30W in combination with Guthion emulsifiable concentrate.
  3. Do not apply OMITE-30W in combination with foliar spray oils.
  4. Do not use additional spray adjuvants.

B. OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

**COMPATIBILITIES —** Except as noted above, OMITE-30W is compatible with most pesticides. (See caution statements.) Check the Omite use booklet or with your supplier for details.

**IMPORTANT NOTICE —** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, **but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.**

2/2