

**PRECAUTIONARY  
STATEMENTS  
HAZARDS TO HUMANS  
DANGER**

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Flammable, keep away from heat and open flame.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**1. PROHIBITIONS**

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

**2. PESTICIDE DISPOSAL**

Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**3. CONTAINER DISPOSAL**

Triple rinse (or equivalent) and offer for recycling, or disposal in approved landfill or bury in a safe place.

**4. GENERAL**

Consult Federal, State or Local Disposal Authorities for approved alternative procedures. Substances prohibited from burning.

**OMITE®-30W**

AGRICULTURAL MITICIDE  
(WEST OF THE ROCKY MOUNTAINS)



**COMPOSITION**

Active Ingredient: (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	30.0%
Inert Ingredients:	70.0%
Total:	100.0%

\*U.S. Patents 3,272,854 - 3,463,859

**KEEP OUT OF REACH OF CHILDREN.  
DANGER**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**IF INHALED:** See a physician if an abnormal reaction occurs.

**IF ON SKIN:** Wash thoroughly with soap and water.

**IF IN EYES:** Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-82-AA

©1969, 1971, 1973 UNIROYAL, Inc. All Rights Reserved.

**NET WEIGHT**

12-14-78

ACCEPTED  
FEB 22 1979  
400-82

**DIRECTIONS FOR USE**

A dry wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two-spotted spider and Willamette mites.

OMITE-30W is not systemic in action; therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-30W per acre. OMITE-30W can be used in concentrate sprays.

**DOSAGE INSTRUCTIONS**

See other side for Dosage Instructions Charts.

400-82

2-22-1979

1/2

**DOSAGE INSTRUCTIONS**

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
ALMONDS (California only)	clover European red two-spotted spider strawberry spider Pacific spider	5-22 lbs.	1-1½ lbs.	500-1500	2	28
APPLES	European red two-spotted spider Pacific spider McDaniel	5-12 lbs.	1-1½ lbs.	500-800	3	7
APRICOTS	Pacific spider two-spotted spider	5-9 lbs.	1½ lbs.	300-600	2	14
CALIFORNIA WALNUTS	European red two-spotted spider	7-22 lbs.	1½ lbs.	500-1500	2	14
FIGS	two-spotted spider Pacific spider strawberry spider	3-6 lbs.	1-1½ lbs.	200-400	2	14
GRAPES	Pacific spider strawberry spider Willamette two-spotted spider	5-9 lbs.	1½ lbs.	350-600	2	21
HOPS	two-spotted spider	5 lbs.	2½ lbs.	200	2	14
NECTARINES	Pacific spider brown almond	5-9 lbs.	1½ lbs.	300-600	2	14
LEMONS (California only)	citrus red	10-15 lbs.	1-1½ lbs.	600-1500	2	7
PEACHES	European red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5-15 lbs.	1½ lbs.	300-1000	2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5-15 lbs.	1½ lbs.	300-1000	2	14
ROSES (greenhouse grown)	two-spotted spider Pacific spider strawberry spider	—	1 lb.	—	—	—

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER CROP SEASON (BLOOM-HARVEST)	LAST SPRAY DAYS BEFORE HARVEST
STRAWBERRIES (See application notes below)	Pacific spider strawberry spider two-spotted spider	5-10 lbs.	5 lbs.	100-200	3*	3

\*Do not apply more than 3 applications of Omite in any 40 day period during the harvesting season.

**APPLICATION NOTES:**

- Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur.
- California: Do not apply at day temperatures above 80°F as fruit burn may occur. (Strawberries only).
- OMITE-30W may cause some leaf burn on new growth (feather growth) of lemons.
- Omite performance is best when daily temperatures average above 70°F.

**POST HARVEST ONLY**

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
NAVEL ORANGES	CITRUS RED	10-15 lbs.	1-1½ lbs.	600-1500	1	See CAUTION BELOW
GRAPEFRUIT	CITRUS RED	10-15 lbs.	1-1½ lbs.	600-1500	1	See CAUTION BELOW

**CAUTION: DO NOT USE ON NAVEL ORANGES OR GRAPEFRUIT AFTER NEW FRUIT IS EXPOSED IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.**

**FOR POST HARVEST AND NON-BEARING TREES ONLY**

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

**DOSAGE INSTRUCTIONS**

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS.	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two-spotted spider Pacific spider McDaniel	6 lbs.	1½ lbs.	400	—	—
PEARS	European red two-spotted spider	5-7½ lbs.	1½ lbs.	300-500	—	—

**USE RESTRICTIONS —** Do not feed livestock on cover crops grown among the trees listed on this label.

**CAUTION:**

- Leaf injury and/or fruit spotting may occur if the following conditions are not observed.
  - Do not apply OMITE-30W with Diazinon-WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
  - OMITE-30W is not compatible with foliar spray oils. (Further information available from supplier or Uniroyal.)
- OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

**COMPATIBILITIES —** Except as noted above, OMITE-30W is compatible with most pesticides. (See caution statements.) Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

**IMPORTANT NOTICE —** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

2/2