

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS DANGER

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited

2. PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a land fill approved for pesticides or buried in a safe place away from water supplies

3. CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling, or disposal in approved land fill or bury in a safe place

4. GENERAL

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning

OMITE®-30W

AGRICULTURAL MITICIDE
(WEST OF THE ROCKY MOUNTAINS)

UNIOYAL

COMPOSITION

Active Ingredient (% by weight)	
Propargite 2-(p-tert-butylphenoxy)- cyclohexyl 2-propynyl sulfite*	30.0%
Inert Ingredients	70.0%
Total	100.0%

*U.S. Patents 3,272,854 - 3,463,859

DIRECTIONS FOR USE

A dry wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two-spotted spider and Willamette mites

OMITE 30W is not systemic in action, therefore complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE 30W per acre. OMITE-30W can be used in concentrate sprays

DOSAGE INSTRUCTIONS

See other side for Dosage Instructions Charts

AL USUARIO:

Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente

TO THE USER:

If you cannot read or understand English, do not use this product until the label has been fully explained to you

KEEP OUT OF REACH OF CHILDREN. DANGER

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water

IF IN EYES: Immediately flush eyes with plenty of water. See a physician

See left side panel for Additional Precautionary Statement

Mfg. by UNIOYAL CHEMICAL, Division of UNIOYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-82-AA

Net Weight

NET WEIGHT

4-18-80

128

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
				Ground	Aerial		
NDS (w/ty)	clover	5.22 lbs	1 1/2 lbs	500-1500	—	2	28
	European red two-spotted spider strawberry spider Pacific spider						
VIA	European red two-spotted spider Pacific spider McDaniel	5.12 lbs	1 1/2 lbs	500-800	—	2	7
	Pacific spider two-spotted spider	5.9 lbs	1 1/2 lbs	700-600	—	2	14
VIA	European red two-spotted spider	7.22 lbs	1 1/2 lbs	500-1500	—	2	14
	two-spotted spider Pacific spider strawberry spider	3-6 lbs	1 1/2 lbs	200-400	—	2	14
	Pacific spider strawberry spider Whamette two-spotted spider	5-9 lbs	1 1/2 lbs	350-600	—	2	21
	two-spotted spider	5 lbs	2 1/2 lbs	200	—	2	14
	Pacific spider brown almond	5-9 lbs *	1 1/2 lbs	700-600	Minimum 20 gal	2	14
	citrus red	10-15 lbs	1 1/2 lbs	600-1500	—	2	7
	European red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5-15 lbs **	1 1/2 lbs	300-1000	Minimum 20 gal	2	14
	European red two-spotted spider Pacific spider	5-15 lbs, **	1 1/2 lbs	300-1000	Minimum 20 gal	2	14
	two-spotted spider Pacific spider strawberry spider	—	1 lb.				
	two-spotted spider Pacific spider	2.5 lbs	2.5 lbs	50-300	—		
	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER CROP SEASON (BLOOM-HARVEST)	LAST SPRAY DAYS BEFORE HARVEST	
	Pacific spider strawberry spider two-spotted spider	5-10 lbs	5 lbs	100-200	3*	3	

are than 3 applications of Omite in any 10 day period during the harvesting

nds of Omite 30W per acre when applied by air
unds of Omite 30W per acre when applied by air

APPLICATION NOTES

- Do not mix with other chemical sprays or liquids when using on strawberries and lemons.
- Do not apply if day temperatures above 80°F as fruit burn may occur (Strawberries only).
- OMITE 30W may cause some leaf burn on new growth (feather growth) of lemons.
- Omite performance is best when daily temperatures average above 70°F.

POST HARVEST ONLY

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
NAVEL ORANGES	CITRUS RED	10-15 lbs	1-1 1/2 lbs	600-1500	1	See CAUTION BELOW
GRAPEFRUIT	CITRUS RED	10-15 lbs	1-1 1/2 lbs	600-1500	1	See CAUTION BELOW

CAUTION: DO NOT USE ON NAVEL ORANGES OR GRAPEFRUIT AFTER NEW FRUIT IS EXPOSED IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	OMITE-30W PER 100 GALLONS OF WATER WEST OF ROCKY MTS	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two-spotted spider Pacific spider McDaniel	6 lbs	1 1/2 lbs	400	—	—
PEARS	European red two-spotted spider	5-7 lbs	1 1/2 lbs	300-500	—	—

USE RESTRICTIONS—Do not feed livestock on cover crops grown among the trees listed on this label

CAUTION:

- Leaf injury and/or fruit spotting may occur if the following conditions are not observed:
 - Do not apply OMITE-30W with Diazinon-WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
 - OMITE-30W is not compatible with foliar spray oils (Further information from supplier or Uniroyal)
- OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

COMPATIBILITIES—Except as noted above, OMITE 30W is compatible with most pesticides. (See caution statements.) Check with your supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.