






IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



- Controls a broad spectrum of mites
- Fits integrated control programs



Revised 1/38

GENERAL INSTRUCTIONS

A dry wettable powder for preparing sprays to control brown almond, citrus, Daniel, Pacific spider, peach silver, strawberry spider, two-spotted spider. OMITE-30W is not systemic in action; therefore, complete and uniform coverage of surfaces and of fruit is necessary for effective control. Always use at least the amount of OMITE-30W per acre. OMITE-30W can be used in concentrate sprays.

AL USUARIO:
Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta sea traducida.

TO THE USER:
If you cannot read or understand English, do not use this product until the label is translated.

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	
			Ground	Aerial
Almonds	European red, two-spotted spider, strawberry spider, Pacific spider	5.22 lbs	500-1000	—
Apples	European red, two-spotted spider, Pacific spider, McDaniel	1.12 lbs	100-200	—
Citrus	Pacific spider, two-spotted spider	1.12 lbs	100-200	—
Plums	Two-spotted spider, Pacific spider, strawberry spider	1.12 lbs	100-200	—
Walnuts	Pacific spider, European red, two-spotted spider, strawberry spider	1.12 lbs	100-200	—

DOSAGE INSTRUCTIONS continued (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
APPLES	Two spotted spider	1 lb.	10-15		2	14
LEMONS LIMES ORANGE	Citrus red	1 lb.	10-15		2	14
NEUTRALS	Pacific spider Strawberry spider	1 lb.	10-15	Medium	2	14
ORNAMENTALS (See application notes)	Two spotted spider	1 lb. per 100 gallons of water	10-15			
PEACHES	European spider Two spotted spider Strawberry spider McDaniel Green spider	1 lb.	10-15	Medium	2	14
PLUMS PRUNES	European spider Two spotted spider Pacific spider	1 lb.	10-15	Medium	2	14
PLUM Greenhouse grown	Two spotted spider Pacific spider Strawberry spider	1 lb. per 100 gallons of water	10-15			
ROSES (field grown)	Two spotted spider Pacific spider	1 lb.	10-15			
STRAWBERRIES (See application notes page 3)	Two spotted spider Strawberry spider Two spotted spider	1 lb.	10-15		2	14
WALNUTS	European spider Two spotted spider	1 lb.	10-15		2	14

† At end of booklet see list of ornamentals which may show signs of phytotoxicity. ‡ Do not apply more than 3 applications of Omite in any 40 day period during the harvesting season. * Use 7 1/2 9 pounds of Omite-30W per acre when applied by air. ** Use 7 1/2 10 pounds of Omite-30W per acre when applied by air.

TECH

APPLICATION NOTES:

1. Complete and uniform coverage is necessary for optimum results.
2. Do not reduce the recommended rate per acre when spraying.
3. Do not combine with other chemicals or spray adjuvants may occur.
4. California: Do not apply at day temperatures above 85°F.
5. OMITE-30W may cause some leaf burn on new growth.
6. Omite performance is best when daily temperatures are above 60°F.
7. Leaf injury and/or fruit spotting may occur if the following conditions exist:
Do not apply OMITE-30W with DIAZINON-WP if a foliar spray of apples and hops.
OMITE-30W is not compatible with foliar spray of Uniroyal.
8. OMITE-30W is not compatible with alkaline materials. The effectiveness of OMITE-30W will be reduced.
9. Ornamentals - 1 lb. of OMITE-30W per 100 gallons of water. The user should initially treat only a small portion of the plant.

POST HARVEST

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT
NAVEL ORANGES	CITRUS RED	10-15 lbs
GRAPEFRUIT	CITRUS RED	10-15 lbs

CAUTION: DO NOT USE ON NAVEL ORANGES & GRAPEFRUIT IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.

of the Rocky Mountains)

GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HAR- VEST
Ground	Aerial		
10	—	2	14
20	—	2	7
30	Minimum 20 gal	2	14
40	—	—	—
50	Minimum 20 gal	2	14
60	—	—	—
70	—	—	—
80	Minimum 20 gal	2	14
90	—	—	—
100	Minimum 20 gal	2	14
110	—	—	—
120	—	—	—
130	—	—	—
140	—	—	—
150	—	—	—
160	—	—	—
170	—	—	—
180	—	—	—
190	—	—	—
200	—	—	—
210	—	—	—
220	—	—	—
230	—	—	—
240	—	—	—
250	—	—	—
260	—	—	—
270	—	—	—
280	—	—	—
290	—	—	—
300	—	—	—
310	—	—	—
320	—	—	—
330	—	—	—
340	—	—	—
350	—	—	—
360	—	—	—
370	—	—	—
380	—	—	—
390	—	—	—
400	—	—	—
410	—	—	—
420	—	—	—
430	—	—	—
440	—	—	—
450	—	—	—
460	—	—	—
470	—	—	—
480	—	—	—
490	—	—	—
500	—	—	—
510	—	—	—
520	—	—	—
530	—	—	—
540	—	—	—
550	—	—	—
560	—	—	—
570	—	—	—
580	—	—	—
590	—	—	—
600	—	—	—
610	—	—	—
620	—	—	—
630	—	—	—
640	—	—	—
650	—	—	—
660	—	—	—
670	—	—	—
680	—	—	—
690	—	—	—
700	—	—	—
710	—	—	—
720	—	—	—
730	—	—	—
740	—	—	—
750	—	—	—
760	—	—	—
770	—	—	—
780	—	—	—
790	—	—	—
800	—	—	—
810	—	—	—
820	—	—	—
830	—	—	—
840	—	—	—
850	—	—	—
860	—	—	—
870	—	—	—
880	—	—	—
890	—	—	—
900	—	—	—
910	—	—	—
920	—	—	—
930	—	—	—
940	—	—	—
950	—	—	—
960	—	—	—
970	—	—	—
980	—	—	—
990	—	—	—
1000	—	—	—

APPLICATION NOTES:

1. Complete and uniform coverage is necessary for optimum results.
2. Do not reduce the recommended rate per acre when applying OMITE-30W in concentrate sprays.
3. Do not combine with other chemicals or spray adjuvants when using on strawberries as leaf and fruit injury may occur.
4. California: Do not apply at day temperatures above 80°F as fruit burn may occur (Strawberries only).
5. OMITE-30W may cause some leaf burn on new growth (feather growth) of lemons.
6. Omite performance is best when daily temperatures average above 70°F.
7. Leaf injury and/or fruit spotting may occur if the following conditions are not observed:
Do not apply OMITE-30W with DIAZINON-VWP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
OMITE-30W is not compatible with foliar spray oils (Further information available from supplier or Uniroyal).
8. OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced.
9. Ornamentals - 1 lb. of OMITE-30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions.

POST HARVEST ONLY

CROP	MITE CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
NAVEL ORANGES	CITRUS RED	10 1/2 oz	600-1500	1	See CAUTION BELOW
GRAPEFRUIT	CITRUS RED	10 1/2 oz	600-1500	1	See CAUTION BELOW

CAUTION: DO NOT USE ON NAVEL ORANGES OR GRAPEFRUIT AFTER NEW FRUIT IS EXPOSED IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.

†† Do not apply more than 3 applications of Omite in a year per acre when applied by air. ** Use 7 1/2 10 pounds

106

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two spotted spider Pacific spider McDaniel	6 lbs	400		
PEARS	European red two spotted spider	5 7/8 lbs	FOR POST HARVEST TREES ONLY 300-500		

USE RESTRICTIONS — Do not feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES — Except as noted above, OMITE-30W is compatible with many pesticides. (See application note 7 (page 3.) Check with your supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

DOSAGE INSTRUCTIONS (EAST OF THE ROCKY MOUNTAINS)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR
APPLES	European red two spotted spider Pacific spider McDaniel	5-12 lbs	400-800	
CARNATIONS CHRYSANTHEMUMS ROSES	two spotted spider	1 1/2 lbs per 100 gallons of water	Spray to wet	
ORNAMENTALS* Field or Nursery	two spotted spider	1 lb per 100 gallons of water	Spray to wet	
PEACHES	European red two spotted spider peach silver strawberry spider McDaniel Pacific spider	5-15 lbs	400-1000	
PEANUTS	two spotted spider	3-5 lbs	See application note 7	
PLUMS PRUNES	European red two spotted spider Pacific spider	5-15 lbs	400-1000	
STRAWBERRIES (Florida and Louisiana only)	two spotted spider	2 1/2	100	(per year)

* At end of booklet see list of ornamentals which may show signs of phytotox.
** Do not apply more than 3 applications of Omite in any 21 day period during

130c

TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
3	1
—	—
—	—
2	14
2	14
2	14
3** (per crop season)	3

5

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
PEARS	European red two spotted spider				

1. Complete and uniform coverage is necessary for optimum results.
2. Do not reduce the recommended rate per acre when applying OMITE 30W in concentrate sprays.
3. Do not apply OMITE 30W under the following conditions as leaf injury and fruit scuffing may occur:
 - when air temperatures exceed 90°F
 - in combination with Guthion emulsifiable concentrate
 - in combination with foliar spray oils
4. Do not use additional spray adjuvants.
5. OMITE 30W is not compatible with a lime materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE 30W will be reduced.
6. Ornamentals: 1 lb. of OMITE 30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions.
7. Peanuts: Premix OMITE 30W with a small amount of water to form a slurry before adding to spray tank. Apply in sufficient water to provide complete coverage of foliage when using low gallonage equipment. A minimum of 20 gallons of water per acre is recommended for best results.
8. Roses: Use is limited to greenhouse, slat and saran grown. Because cultivar reaction and cultural practices differ widely, the user should initially treat only a small portion of his crop to confirm plant safety under his conditions. Repeat applications of OMITE 30W are normally necessary for extended control.

6

[illegible]

use that will not bear fruit within one year of Omite Application

MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
European red Two spotted spider	5.7 55	400 50		

emulsifiable concentrate
or spray oils

1.5-3.0W per 100 gallons of water equals 1 ounce (28.4 grams) per 625 gallons. Only a small portion of his crop to confirm plant safety under his conditions. Mix with a small amount of water to form a slurry before adding to spray tank, provide complete coverage of foliage when using low gallonage equipment. A 1/2 gallon per acre is recommended for best results.

of 0.05 and 0.01 are normally necessary for extended work

COMPATIBILITIES — Except as noted above, OMITE-30W is compatible with most pesticides. (See application notes 3 & 4) Check the Omite use booklet or with your supplier for details.

Plants Showing Slight Necrosis (leaf burn)

[illegible]

Major's name is Major Steele
Philadelphia, Pennsylvania
He is a Major in the
United States Army
He is a Major in the
United States Army
He is a Major in the
United States Army