


BEST DOCUMENT AVAILABLE

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

RECEIVED

SEP 02 1982

Registration Division (TS-767)



- Controls a broad spectrum of mites
- Fits integrated control programs

SEP 02 1982

Registration Division (TS-767)

11-10-81

GENERAL INSTRUCTIONS

A dry, wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two spotted spider and Williamette mites. OMITE-30W is not systemic in action, therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-30W per acre. OMITE-30W can be used in concentrate sprays.

AL USUARIO

Se utiliza en polvo para preparar pulverizaciones para controlar ácaros: almendra roja, rojo de cítricos, rojo de alfalfa, rojo de Europa, rojo de McDaniel, araña del Pacífico, plata de melocotón, araña de la fresa, araña de dos manchas y araña de Williamette.

TO THE USER

If you cannot read or understand English, do not use this product until the label has been fully explained to you.

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Air		
ALMOND	Brown almond mite	1.0	1.0	1.0	1	14
APPLE	European red mite, Pacific spider mite, two spotted spider mite, Williamette mite	1.0	1.0	1.0	1	14
CHERRY	European red mite, Pacific spider mite, two spotted spider mite, Williamette mite	1.0	1.0	1.0	1	14
CITRUS	Citrus red mite	1.0	1.0	1.0	1	14
CLOVER	Clover mite	1.0	1.0	1.0	1	14
PEACH	Peach silver mite	1.0	1.0	1.0	1	14
STRAWBERRY	Strawberry spider mite	1.0	1.0	1.0	1	14
WILLAMETTE	Williamette mite	1.0	1.0	1.0	1	14

DOSAGE INSTRUCTIONS continued (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
HOPS	two spotted spider	5 lbs	200	—	2	14
LEMONS (CA & AZ)	citrus red	7.5 lbs	2500	—	2	7
NECTARINES	Pacific spider brown almond	5.9 lbs *	500-600	Minimum 20 gal	2	14
ORNAMENTALS (field & nursery)	two spotted spider	1 lb per 100 gallons of water	Spray to wet	—	—	—
PEACHES	European red Pacific spider two spotted spider strawberry spider McDaniel peach silver	5.15 lbs **	300-1000	Minimum 20 gal	2	14
PLUMS PRUNES	European red two spotted spider Pacific spider	5.15 lbs. **	300-1000	Minimum 20 gal	2	14
ROSES (greenhouse grown)	two spotted spider Pacific spider strawberry spider	1 lb per 100 gallons of water	—	—	—	—
ROSES (field grown)	two spotted spider Pacific spider	2.5 lbs.	50-300	—	—	—
STRAWBERRIES (See application notes page 3)	Pacific spider strawberry spider two spotted spider	5.10 lbs	500-200	—	3†† (per crop season)	3
WALNUTS (CA only)	European red two spotted spider	7.22 lbs	500-1500	—	2	14

† At end of booklet, see list of ornamentals which may show signs of phytotoxicity. †† Do not apply more than 3 applications of Omite in any 40 day period during the harvesting season. * Use 7½-9 pounds of Omite-30W per acre when applied by air. ** Use 7½-10 pounds of Omite-30W per acre when applied by air.

AT DOCUMENT AVAILABLE

APPLICATION NOTES:

- 1 Complete and uniform coverage is necessary for optimum results
- 2 Do not reduce the recommended rate per acre when applying OMITE-30W in concentrate sprays
- 3 Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur
- 4 California: Do not apply at day temperatures above 80°F as fruit burn may occur (Strawberries only).
- 5 OMITE 30W may cause some leaf burn on new growth (leather growth) of lemons
- 6 Omite performance is best when daily temperatures average above 70°F
- 7 Leaf injury and/or fruit spotting may occur, if the following conditions are not observed
 - Do not apply OMITE 30W with DIAZINON—WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops
 - OMITE-30W is not compatible with foliar spray oils (Further information available from supplier or Uniroyal)
- 8 OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced
- 9 Ornamentals - 1lb of OMITE-30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions
- 10 Keep workers out for seven (7) days after application. There is a 7-day safety reentry interval for grapes. Workers shall not enter treated areas to perform activities that involve substantial or prolonged contact with treated foliage until the 7-day safety reentry has expired

POST HARVEST ONLY

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
NAVEL ORANGES	CITRUS RED	10-15 lbs	8-10-1500	1	See CAUTION BELOW
GRAPEFRUIT	CITRUS RED	10-15 lbs	8-10-1500	1	See CAUTION BELOW

CAUTION: DO NOT USE ON NAVEL ORANGES OR GRAPEFRUIT AFTER NEW FRUIT IS EXPOSED IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two spotted spider Pacific spider McDaniel	6 lbs	400		
PEARS	European red two spotted spider	5 7½ lbs	FOR POST HARVEST TREES ONLY 300-500		

USE RESTRICTIONS — Do not feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES — Except as noted above, OMITE 30W is compatible with many pesticides. (See application note 7 (page 3) Check with your supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

BEST DOCUMENT AVAILABLE

DOSAGE INSTRUCTIONS (EAST OF THE ROCKY MOUNTAINS)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
APPLES	European red two spotted spider Pacific spider McDaniel	5.12 lbs	400-800	3	7
CARNATIONS CHRYSANTHEMUMS ROSES	two spotted spider	1 1/2 lbs. per 100 gallons of water	Spray to wet	---	---
ORNAMENTALS* Field or Nursery	two spotted spider	1 lb. per 100 gallons of water	Spray to wet	---	---
PEACHES	European red two spotted spider Pacific spider McDaniel	5.15 lbs	400-1000	2	14
PEANUTS	two spotted spider	3.5 lbs	See application note 7	2	14
PLUMS PRUNES	European red two spotted spider Pacific spider	5.15 lbs	400-1000	2	14
STRAWBERRIES (Florida and Louisiana only)	two spotted spider	2 1/2	100	3** (per crop season)	3

- * All types of bark or seed coat may show signs of phytotoxicity.
 ** Do not apply more than 3 applications of Omite in any 21 day period during the harvesting season.

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

DOSAGE INSTRUCTIONS (EAST OF THE ROCKY MOUNTAINS)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE PER TREATMENT	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
PEANUTS	European red two spotted spider	5.7-11.4 lbs	40-80 gal		

APPLICATION NOTES:

1. Complete and uniform coverage is necessary for optimum results.
2. Do not reduce the recommended rate per acre when applying OMITE-30W in concentrate sprays.
3. Do not apply OMITE-30W under the following conditions as leaf injury and fruit spotting may occur:
 - when air temperatures exceed 90°F
 - in combination with Guthion emulsifiable concentrate
 - in combination with foliar spray oils
4. Do not use additional spray adjuvants.
5. OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced.
6. Ornamentals - 1 lb. of OMITE-30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.5 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions.
7. Peanuts - Premix OMITE-30W with a small amount of water to form a slurry before adding to spray tank. Apply in sufficient water to provide complete coverage of foliage when using low gallonage equipment. A minimum of 20 gallons of water per acre is recommended for best results.
8. Roses - Use is limited to greenhouse, slat and saran grown. Because cultivar reaction and cultural practices differ widely, the user should initially treat only a small portion of his crop to confirm plant safety under his conditions. Repeat applications of OMITE-30W are normally necessary for extended control.

6

BEST DOCUMENT AVAILABLE

USE RESTRICTIONS — Do not graze or feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES — Except as noted above, OMITE-30W is compatible with most pesticides. (See application notes 3 & 4). Check the Omite use booklet or with your supplier for details.

The following ornamentals may show some sign of phytotoxicity.

Plants Showing Slight Necrosis (leaf burn)

<i>Abelia grandiflora</i>	<i>Cotoneaster</i> "Skogsholmen"	<i>Hydrangea macrophylla</i> "Merrill's Beauty"
<i>Acer palmatum</i>	<i>Cupania anacardioides</i>	<i>Hydrangea macrophylla</i> "Revelation"
<i>Alectryon subcuneatus</i>	<i>Eucalyptus cinerea</i>	<i>Hydrangea macrophylla</i> "Trophy"
<i>Aralia setoides</i>	<i>Eucalyptus globulus compacta</i>	<i>Hydrangea macrophylla</i> "White"
<i> Aucubia japonica picturata</i>	<i>Eucalyptus niphophila</i>	<i>Ilex</i> "Nellie Stevens"
<i>Aucubia japonica variegata</i>	<i>Eucalyptus rostrata</i>	<i>Koeleria paniculata</i>
<i>Beaumontia grandiflora</i>	<i>Eucalyptus sideroxylon rosea</i>	<i>Lagerstroemia</i> "Petite Orchid"
<i>Berberis thunbergii atropurpurea</i>	<i>Eucalyptus teretifolia</i>	<i>Lagerstroemia</i> "Petite Pinkie"
<i>Berberis thunbergii atropurpurea nana</i>	<i>Eucalyptus viminalis</i>	<i>Mahonia repens</i>
<i>Crimson Pigeon</i>	<i>Gardenia jasminoides</i> "Mystery"	<i>Nyssa sylvatica</i>
<i>Bauhinia variegata</i>	<i>Gardenia jasminoides</i> "Vendula"	<i>Pachysandra terminalis</i>
<i>Buxus microphylla</i> "Richard"	<i>Hedera canariensis</i>	<i>Philadelphus</i> "Lance"
<i>Cissus mandariniana</i>	<i>Hedera</i> "Green Tree"	<i>Prunus integrifolia</i>
<i>Cornus latifolia</i>	<i>Hydrangea macrophylla</i> "Hamburg"	<i>Rosa Banksiae alba</i>
<i>Cotoneaster horizontalis perposita</i>	<i>Hydrangea macrophylla hortensis compacta</i>	<i>Schinus terebinthifolius</i>
<i>Cotoneaster horizontalis repindens</i>	<i>Hydrangea macrophylla hortensis</i> "Excelsior"	<i>Taxus canadensis</i>
		<i>Weigela</i> "Brilliant Snowflake"
		<i>Podocarpus gracilis</i>

Plants Showing Moderate to Severe Necrosis (leaf burn)

<i>Aralia nudicaulis</i>	<i>Cotoneaster</i> "Lata Formosa"	<i>Magnolia grandiflora</i> "Majestic Beauty"
<i>Amelanchier canadensis</i>	<i>Euphorbia corollata</i>	<i>Philadelphus</i> "Minnesota Snowflake"
<i>Aucubia japonica</i> "Dwarf Female"	<i>Eucalyptus laetifolia</i>	<i>Philadelphus virginicus</i>
<i>Beloperone longifolia</i>	<i>Geum sibiricum</i>	<i>Potentilla fruticosa</i> "Moonlight"
<i>Berberis</i> "Mentorena"	<i>Hamamelis</i>	<i>Potentilla grandiflora</i>
<i>Ceanothus</i> "Ray Hartman"	<i>Hedera</i> "Hera"	<i>Quercus alba</i>
<i>Cissus repens</i>	<i>Hydrangea macrophylla</i>	<i>Taxus canadensis</i>
<i>Cissus hypoglauca</i>	<i>Lythrum hyssopifolium</i>	<i>Taxus canadensis</i>



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

310/80757
17/19

JAN 3 1983

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

Uniroyal Chemical
Crop Protection, Research & Devel.
74 Amity Rd.
Bethany, CT 06525

Attn: Mr. Bogislav von Schmeling

Dear Mr. von Schmeling:

Subject: Amendment - Arizona to Almonds
Omite 30W
EPA Reg. No: 400-82
Omite 6E
EPA Reg. 400-89
Appl. Dates August 25, 1982

The labeling for the product registration referred to above, submitted in connection with amended registration, under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, subject to the following comments, and stamped copies are enclosed for your records.

1. For the Omite 30W we received only the application instructions booklet; please make the same revision on the container label.
2. At the next label printing, for both products, from the "If swallowed" STATEMENT OF PRACTICAL TREATMENT, delete "Call a physician immediately." and insert as the first line of the "If swallowed" statement:

"Call a physician or Poison Control Center immediately."

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (15-7b7C)