

SEE FOLDER FOR USE INSTRUCTIONS — ENCLOSED 'N BOTTOM FOLD

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
DANGER**

Corrosive, causes eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS
Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

3. CONTAINER DISPOSAL

Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

4. GENERAL

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

OMITE®-30W

AGRICULTURAL MITICIDE

UNIROYAL

COMPOSITION

Active Ingredient (% by weight)	
Propargite 2-(p-tert-butylphenoxy)- cyclohexyl 2-propynyl sulfite*	30.0%
Inert Ingredients	70.0%
Total	100.0%

*U.S. Patents 3,272,854 - 3,463,859

APR 15 1983

OMITE-30W is a pesticide
registered under
EPA Reg. No. 400-82

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

AL USUARIO:

Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TO THE USER:

If you cannot read or understand English, do not use this product until the label has been fully explained to you.

**KEEP OUT OF REACH OF CHILDREN.
DANGER**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.
Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-82-AA

© 1969 1971 UNIROYAL, Inc. All Rights Reserved

NET WEIGHT

1-21-83

APR 25 1983

EPA Reg. No. 400-82

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

PACKAGE CONTAINS OMITE-30W IN A WATER SOLUBLE BAG-ENTIRE CONTENTS DISSOLVE IN WATER. After opening package, immediately dump entire contents into spray tank. Do not touch bag; handling may cause breakage.
THIS BAG CONTAINS 2x2½ POUND WATER SOLUBLE BAGS.

HOW TO APPLY

OMITE®-30W
AGRICULTURAL MITICIDE
(West of the Rocky Mountains)

UNIROYAL

EPA Reg. No. 100-82-AA

- Controls a broad spectrum of mites
- Fits integrated control programs.

Mfg. by UNIROYAL CHEMICAL
Division of UNIROYAL, Inc.
Naugatuck, Connecticut 06770
©1969, 1971, 1973 UNIROYAL, Inc.
All Rights Reserved

1-18-83

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	Number of 2½ lb. water soluble bags per acre per treatment	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
ALMONDS (CA & AZ)	clover European red two-spotted spider strawberry spider Pacific spider	2-9	500-1500	—	2	28
APPLES	European red two-spotted spider Pacific spider McDaniel	2-5	500-800	—	3	7
APRICOTS	Pacific spider two-spotted spider	2-4	300-600	—	2	14
FIGS	two spotted spider Pacific spider strawberry spider	1-2	200-400	—	2	14
GRAPES 7-Day Reentry Interval (see note 10)	Pacific spider strawberry spider Willamette two spotted spider	2-4	350-600	—	2	21
HOPS	two-spotted spider	2	200	—	2	14
LEMONS (CA & AZ)	citrus red	4-6	600-1500	—	2	7
NECTARINES	Pacific spider brown almond	2-4*	300-600	Minimum 20 gal	2	14
ORNAMENTALS (field & nursery)	two spotted spider	See footnote 9	Spray to wet	—		
PEACHES	European red Pacific spider two spotted spider strawberry spider McDaniel peach silver	2-6**	300-1000	Minimum 20 gal	2	14

POST HARVEST ONLY

CROP	MITES CONTROLLED	Number of 2½ lb. water soluble bags per acre per treatment	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
NAVEL ORANGES	CITRUS RED	4-6	600-1500	1	See CAUTION BELOW
GRAPEFRUIT	CITRUS RED	4-6	600-1500	1	See CAUTION BELOW

CAUTION: DO NOT USE ON NAVEL ORANGES OR GRAPEFRUIT AFTER NEW FRUIT IS EXPOSED IN THE SPRING AS FRUIT SPOTTING MAY OCCUR.

FOR POST HARVEST AND NON-BEARING TREES ONLY

Non-bearing trees are those that will not bear fruit within one year of Omite Application.

DOSAGE INSTRUCTIONS (West of the Rocky Mountains)

CROP	MITES CONTROLLED	Number of 2½ lb. water soluble bags per acre per treatment	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two-spotted spider Pacific spider McDaniel	2	400	—	—
PEARS	European red two-spotted spider	2-3	FOR POST HARVEST TREES ONLY 300 500		

USE RESTRICTIONS — Do not feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES — Except as noted above, OMITE-30W is compatible with many pesticides. (See application note 7 (page 2). Check with your supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

April 20 1963

400-82

DOSAGE INSTRUCTIONS continue (West of the Rocky Mountains)

CROP	MITES CONTROLLED	Number of 2 lb. water soluble bag per acre per treatment	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
PLUMS PRUNES	European red two spotted spider Pacific spider	2-6**	300-1000	Minimum 1-20 gal	2	14
ROSES (greenhouse grown)	two spotted spider Pacific spider strawberry spider	1 per 250 gallons of water	—	—	—	—
ROSES (field grown)	two-spotted spider Pacific spider	1-2	50-300	—	—	—
STRAWBERRIES (See application notes)	Pacific spider strawberry spider two-spotted spider	2-4	100-200	—	3†† (per crop season)	3
WALNUTS (CA only)	European red two spotted spider	3-9	500-1500	—	2	14

† At end of booklet, see list of ornamentals which may show signs of phytotoxicity. †† Do not apply more than 3 applications of Omite in any 40 day period during the harvesting season. * Use 7 1/2-9 pounds of Omite-30W per acre when applied by air. ** Use 7 1/2-10 pounds of Omite-30W per acre when applied by air.

APPLICATION NOTES:

1. Complete and uniform coverage is necessary for optimum results.
2. Do not reduce the recommended rate per acre when applying OMITE-30W in concentrate sprays.
3. Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur.
4. California: Do not apply at day temperatures above 80°F as fruit burn may occur. (Strawberries only).
5. OMITE-30W may cause some leaf burn on new growth (feather growth) of lemons.
6. Omite performance is best when daily temperatures average above 70°F.
7. Leaf injury and/or fruit spotting may occur, if the following conditions are not observed:
 - Do not apply OMITE-30W with DIAZINON-WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
 - OMITE-30W is not compatible with foliar spray oils. (Further information available from supplier of Uniroyal).
8. OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced.
9. Ornamentals - 2 1/2 lb. of OMITE-30W per 250 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions.
10. Keep workers out for seven (7) days after application. There is a 7-day safety reentry interval for grapes. Workers shall not enter treated areas to perform activities that involve substantial or prolonged contact with treated foliage until the 7-day safety reentry has expired.

The following ornamentals may show some sign of phytotoxicity.

Plants Showing Light Necrosis (leaf burn)

Abelia grandiflora	Gardenia jasminoides 'Vitchi'
Acer palmatum	Hedera canariensis
Alectryon subcinerus	Hedera 'Ghost Tree'
Aralia seiboldi	Hydrangea macrophylla 'Hamburg'
Aucuba japonica picta	Hydrangea macrophylla hortensis compacta
Aucuba japonica variegata	Hydrangea macrophylla hortensis
Beaumontia grandiflora	Excelsior
Bergia thunbergia atropurpurea	Hydrangea macrophylla 'Merritt's Beauty'
Bergia thunbergia atropurpurea f. nana	Hydrangea macrophylla 'Revelation'
Crimson Pygmy	Hydrangea macrophylla 'Trophee'
Bougainvillea 'Reddwarf'	Hydrangea macrophylla 'White'
Buxus microphylla 'Richard'	Illex 'Nellie Stevens'
Cissus mandarin	Koeleria paniculata
Coccoloba laurifolia	Lagerstroemia 'Petite Orchid'
Cotoneaster 'Mormon'	Lagerstroemia 'Petite Pinkie'
Cotoneaster 'Sagittaria'	Malva repens
Cotoneaster 'Skogholmen'	Nyssa sylvatica
Cupania anacardioides	Pachysandra terminalis
Eucalyptus cinerea	Philadelphus 'Glacier'
Eucalyptus globus compacta	Peccaropus gracilior
Eucalyptus niphophila	Prunus integrifolia
Eucalyptus rostrata	Rosa Banksiae alba
Eucalyptus sideroxylon rosea	Schinus terebinthifolius
Eucalyptus torquata	Tetonia stans
Eucalyptus viminalis	Weigela 'Bristol Snowflake'
Gardenia jasminoides 'Mystery'	

Plants Showing Moderate to Severe Necrosis (leaf burn)

Acacia podalyriaefolia
 Alnus rhombifolia
 Aucuba japonica Dwarf Female
 Beloperone tomentosa
 Berbis Mentorensis
 Ceanothus Ray Hartman
 Cissus capensis
 Cissus hypoglauca
 Cotoneaster bullata floribunda
 Dendromecon harfordii thamnoides
 Eucalyptus ficifolia
 Gelsemium sempervirens
 Grevillea Noelli

Hedera helix Goldheart
 Leptospermum scoparium Keatleyi
 Liquidambar orientalis
 Magnolia grandiflora Majestic Beauty
 Philadelphus virginialis
 Potentilla fruticosa Moonlight
 Philadelphus Minnesota Snowflake
 Potentilla fruticosa Moonlight
 Potentilla grandiflora
 Quercus agrifolia
 Ternstroemia japonica
 Tristania conferta

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
DANGER**

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

This is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide spray, mixture or residue that cannot be used, or chemically reprocessed should be disposed of in a land fill approved for pesticides or buried in a safe place away from water supplies.

3. CONTAINER DISPOSAL

Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

4. GENERAL

Consult Federal, State and Local Disposal Authorities for approved alternative procedures, such as incineration or open burning.

OMITE[®]-CR
AGRICULTURAL MITICIDE



COMPOSITION

Active Ingredients (1% by weight)	
Propargite 2-(p-tert-butyl phenoxy)-	30.0%
cyclohexyl 2-propynyl sulfite*	70.0%
Inert Ingredients	
Total	100.0%

*U.S. Patents 3,272,854 - 3,463,859

**KEEP OUT OF REACH OF CHILDREN.
DANGER**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

IF INHALED See a physician if an abnormal reaction occurs.

IF ON SKIN Wash thoroughly with soap and water.

IF IN EYES Immediately flush eyes with plenty of water. See a physician.

AVOID EXPOSURE TO DUST

See left panel for Additional Precautionary Statement

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-82-AA

1982-1991 © UNIROYAL, Inc. All Rights Reserved

NET WEIGHT

1-19-83

DIRECTIONS FOR USE

OMITE-CR is a dry wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two-spotted spider and Willamette mites.

OMITE-CR is not systemic in action; therefore, complete and uniform coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-CR per acre. OMITE-CR can be used in concentrate sprays.

DOSAGE INSTRUCTIONS

See other side for Dosage Instructions.

MIXING INSTRUCTIONS

This package contains 2 x 1 1/2 pound water soluble bags of OMITE-CR. After opening package:

1. Do not open or perforate the inner water soluble bags. The water soluble bags and contents will dissolve in water.
2. Immediately dump entire contents into spray tank.
3. Do not touch water soluble bags as handling may cause breakage.
4. If only one water soluble bag is used, store the remaining bag in the original package in a dry location away from heat or cold.

AL USUARIO:

Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TO THE USER

If you cannot read or understand English, do not use this product until the label has been fully explained to you.

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	NUMBER OF 1½ POUND WATER SOLUBLE BAGS PER ACRE	GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
ALMONDS (CA only)	European red two spotted spider strawberry spider Pacific spider	3-15	500-1500	—	2	26
APPLES	European red two spotted spider Pacific spider McDaniel	3-8	500-800	—	3	7
APRICOTS	Pacific spider two spotted spider	3-6	300-600	—	2	14
FIGS	two spotted spider Pacific spider strawberry spider	2-3	200-400	—	2	14
GRAPES 7 Day Reentry Interval (See note 6)	Pacific spider strawberry spider Vernette two spotted spider	3-6	350-600	—	2	21
HOPS	two spotted spider	3	200	—	2	14
LEMONS (AZ)	citrus red	7-10	600-1500	—	2	7
NECTARINES	Pacific spider brown almond	3-6*	300-600	Minimum 20 gal	2	14
ORNAMENTALS (ec & nursery)	two spotted spider	1 See footnote 5	Spray to wet	—	—	—
ORANGES GRAPE FRUIT LEMONS (CA only)	citrus red two spotted spider	7-10	1000	—	2	7
PEARS	European red*** two spotted spider McDaniel	3-4	500	—	2	14
PEACHES	European red Pacific spider two spotted spider strawberry spider McDaniel beach silver	3-10**	300-1000	Minimum 20 gal	2	14
PLUMS PRUNES	European red two spotted spider Pacific spider	3-10**	300-1000	Minimum 20 gal	2	14
ROSES (greenhouse grown)	two spotted spider Pacific spider strawberry spider	1 See footnote 5	150	—	—	—
ROSES (field grown)	two spotted spider Pacific spider	1-3 See footnote 5	50-300	—	—	—
STRAWBERRIES (See application notes below)	Pacific spider strawberry spider two spotted spider	3-7	100-200	—	3†	3
WALNUTS (CA only)	European red two spotted spider	5-15	500-1500	—	2	14

APPLICATION NOTES:

- Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur.
- California: Do not apply at day temperatures above 80°F as fruit burn may occur (Strawberries only).
- OMITE-CR may cause some leaf burn on new growth (feather growth) of lemons.
- Omite performance is best when daily temperatures average above 70°F.
- ORNAMENTALS - 1½ lb OMITE-CR per 150 gallons of water equals 1-ounce (28.4 grams) per 6.25 gallons. The user should initially treat only a small portion of his crop to confirm plant safety under his conditions.
- Keep workers out for seven (7) days after application. There is a seven day reentry interval for grapes. Workers shall not enter treated areas to perform activities that involve substantial or prolonged contact with treated foliage until the 7-day reentry interval has expired.
- OMITE-CR when used alone may follow foliar oils at an interval of greater than 40 days after an oil application.
- Foliar oil applications must not follow OMITE-CR treatments within 30 days.

FOR POST HARVEST AND NON-BEARING TREES

Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	NUMBER OF 1½ POUND WATER SOLUBLE BAGS PER ACRE	GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
CHERRIES	European red two spotted spider Pacific spider McDaniel	4	400	—	—

USE RESTRICTIONS—Do not feed livestock on cover crops grown among the trees listed on this label.

CAUTION:

- A. Leaf injury and/or fruit spotting may occur if the following conditions are not observed.
- Do not apply OMITE-CR with Diazinon-WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
 - OMITE-CR is not compatible with foliar spray oils. (Further information from supplier or Uniroyal.)
- B. OMITE-CR is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

COMPATIBILITIES—Except as noted above, OMITE-CR is compatible with most pesticides. (See caution statements.) Check with your supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

† Do not apply more than 3 applications of OMITE-CR in any 40 day period during the harvest season.

* Use 5-6 water soluble bags per acre when applied by air.

** Use 5-7 water soluble bags per acre when applied by air.

*** Requires 5 water soluble bags per acre