



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 12 1987

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

Raymond A. Cardona, Ph.D.
Uniroyal Chemical Company, Inc.
74 Amity Road
Bethany, CT 06525

300, 182 791
17 25

Dear Dr. Cardona:

Subject: Amendment - Label Amendment - Combine East &
West Label Information and Reduce Amount
of Water in Spray Gallonage
Omite 30W
EPA Reg. No. 400-82
Your Application Dated September 9, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. The minimum spray gallonage for apples should be 500 i.e. 500-800.
2. Delete the sentence, "For application at water volumes less than indicated on the chart above xxxxx."


Although the time period for submission of revised labeling in accordance with the Registration Standard on Pro-pargite issued September 30, 1986 has not elapsed we encourage you to incorporate the applicable end-use label revisions recommended under this standard on the subject product labeling.

Submit five (5) copies of your final printed labeling before your release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS**

DANGER

Corrosive, causes eye damage. Harmful if swallowed. Inhaled or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

OMITE®-30W

AGRICULTURAL MITICIDE

ACCEPTED
with COMMENTS
in EPA Letter Dated:

JAN 12 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 400-82

UNIROYAL

COMPOSITION

Active Ingredients: (% by weight)

Propargite [2-(p-tert-butylphenoxy)
cyclohexyl 2-propynyl sulfite]* 30.0%

Inert Ingredients: 70.0%

Total: 100.0%

*U.S. Patent Nos. 3,272,854; 3,463,859

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for additional precautionary statement.

UNIROYAL CHEMICAL — Division of UNIROYAL, Inc.
Naugatuck, Connecticut 06770

EPA REG. NO. 400-82

EPA EST. NO.

OMITE is a Registered Trademark of UNIROYAL, Inc.

©Copyright 1984 Uniroyal, Inc.

5-26-84

Lot. No.

NET WEIGHT:

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to the workers who are expected to be in a treated area or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER, Area to be treated with Propoxite on (date of application). Do not enter without appropriate protective clothing until sprays have dried." Or in the case of grape vineyards, "Do not enter treated areas to perform activities that involve substantial or prolonged contact with treated foliage for 7 days. If accidentally exposed, consult the Statement of Practical Treatment portion of this label for first aid recommendations."

GENERAL INSTRUCTIONS

OMITE-30W is a dry wettable powder for preparing sprays to control *brown mites*, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, strawberry spider, two-spotted spider and Willamette mites.

OMITE-30W is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-30W per acre. OMITE-30W can be used in concentrate sprays, however, do not reduce the recommended rate per acre.

OMITE-30W performance is best when day temperatures average above 70°F.

PRECAUCION AL USUARIO:

Si usted no puede leer ingles no use este producto hasta que la etiqueta haya sido explicada ampliamente.

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	LAST SPRAY DAYS BEFORE HARVEST
			Ground	Aerial		
ALMONDS (CA & AZ)	clover European red two-spotted spider strawberry spider Pacific spider	5-15/62	50	—	2	28
APPLES ¹	European red two-spotted spider Pacific spider McDaniel	5-12 lbs.	40-800	—	3	7
APRICOTS (West of Rocky Mts.)	Pacific spider two-spotted spider	5-9 lbs.	50	—	2	14
CARNATIONS CHRYSANTHEMUMS	two-spotted spider	1-1½ lbs. per 100 gals.	Spray to wet	—	—	—
FIGS	two-spotted spider Pacific spider strawberry spider	3-6 lbs.	50	—	2	14
GRAPES ² 7-Day reentry Interval (West of Rocky Mts.)	Pacific spider strawberry spider Willamette two-spotted spider	5-9 lbs.	40	—	2	21
HOPS ¹	two-spotted spider	5 lbs.	100	—	2	14
LEMONS ¹ (CA & AZ)	citrus red	10-15 lbs.	100	—	2	7
NECTARINES (West of Rocky Mts.)	Pacific spider brown almond	5-9 lbs.*	50	20 gal.	2	14
ORNAMENTALS ¹ (field and nursery)	two-spotted spider	1 lb. per 100 gallons of water	Spray to wet	—	—	—
PEACHES	European red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5-15 lbs.**	50	20 gal.	2	14
PEANUTS	two-spotted spider	3-5 lbs.	20	—	2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5-15 lbs.**	50	20 gal.	2	14
ROSES ¹ (greenhouse. grown)	two-spotted spider Pacific spider strawberry spider	1 lb. per 100 gallons of water	Spray to wet	—	—	—
ROSES ¹ (field grown)	two-spotted spider Pacific spider	2-5 lbs.	50	—	—	—
STRAWBERRIES ¹ ⁵ (West of Rocky Mts.)	Pacific spider strawberry spider two-spotted spider	5-10 lbs.	100	—	3 per 40 day harvest season	3
STRAWBERRIES ¹ ⁵ (LA & FL)	Pacific spider strawberry spider two-spotted spider	2½ lbs.	100	—	3 per 21 day harvest season	3
WALNUTS (CA only)	European red two-spotted spider	7-22 lbs.	100	10	2	14

*Use 7½-9 pounds of OMITE-30W per acre when applied by air. **Use 7½-10 pounds of OMITE-30W per acre when applied by air.

For application at water volumes less than indicated on the chart above, use only equipment designated specifically for high concentrate application. For both dilute and concentrate sprays, always apply sufficient water to obtain thorough, uniform coverage.

Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR
NAVEL ORANGES ¹ (West of Rocky Mts.)	citrus red	10-15 lbs.	100	1
GRAPEFRUIT ¹ (West of Rocky Mts.)	citrus red	10-15 lbs.	100	1
CHERRIES (West of Rocky Mts.)	European red two-spotted spider Pacific spider McDaniel	6 lbs.	50	—
PEARS	European red two-spotted spider	5-7½ lbs.	50	For use after harvest only.

APPLICATION NOTES:

- Apples and Hops: Leaf injury and/or fruit spotting may occur if the following conditions are not observed:
Do not apply OMITE-30W with Diazinon WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.
Do not apply in combination with Guthion emulsifiable concentrate.
Do not apply OMITE-30W with foliar spray oils because of incompatibility. See Compatibilities or contact your supplier or Uniroyal for further information.
- Grapes: When applying OMITE-30W on grapes keep workers out of treated areas) for seven days after application. Workers shall not enter treated areas to perform activities that involve substantial or prolonged contact with treated foliage until the 7 day safety reentry interval has expired (see reentry statement).
- Citrus: OMITE-30W may cause some leaf burn on new growth (leather growth) of lemons.
Foliar oil applications must not follow OMITE-30W treatments within 30 days.
OMITE-30W when used alone may follow foliar oils at an interval of greater than 40 days after an oil application.
Do not use on navel oranges or grapefruit after new fruit is exposed in the spring as fruit spotting may occur.
- Ornamentals and Roses: 1 lb. of OMITE-30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. Because cultivars and cultural practices differ widely, the user should initially treat only a small portion of his crop to confirm plant safety under his conditions. See list of ornamentals on this label which may show phytotoxicity. Repeat applications of OMITE-30W are normally necessary for extended control.
- Strawberries: Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur.
California: Do not apply at day temperatures above 80°F as fruit burn may occur.

COMPATIBILITIES — Except as noted OMITE-30W is compatible with many pesticides (See application notes). Check with your Supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

- Do not use additional spray adjuvants.
- OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced.
- Do not apply in combination with foliar spray oils as phytotoxicity may result.

USE RESTRICTIONS — Do not feed livestock on cover crops grown among the trees listed on this label.

The following ornamentals may show some signs of phytotoxicity.

PLANTS SHOWING SLIGHT NECROSIS (leaf burn)

Abelia grandiflora	Cotoneaster "Skogsholmen"	Hydrangea macrophylla "Merritt's Beauty"
Acer palmatum	Cupania anacardioides	Hydrangea macrophylla "Revelation"
Alectryon subcinerus	Eucalyptus cinerea	Hydrangea macrophylla "Trophee"
Aralia sieboldi	Eucalyptus globulus compacta	Hydrangea macrophylla "White"
Aucuba japonica picturata	Eucalyptus niphophila	Ilex "Nellie Stevens"
Aucuba japonica variegata	Eucalyptus rostrata	Koeleria paniculata
Beaumontia grandiflora	Eucalyptus sideroxylon rosea	Lagerstroemia "Petite Orchid"
Berberis thunbergii atropurpurea	Eucalyptus torquata	Lagerstroemia "Petite Pinkie"
Berberis thunbergii atropurpurea nana	Eucalyptus viminalis	Mahonia repens
"Crimson Pygmy"	Gardenia jasminoides "Mystery"	Nyssa sylvatica
Bougainvillea rhodamine	Gardenia jasminoides "Veilch"	Pachysandra terminalis
Buxus microphylla "Richard"	Hedera canariensis	Philadelphus "Glacier"
Cissus mandarinum	Hedera "Ghost Tree"	Podocarpus gracilior
Coccolus laurifolius	Hydrangea macrophylla "Hamburg"	Prunus integrifolia
Cotoneaster horizontalis	Hydrangea macrophylla hortensis	Rosa Banksiae alba
Cotoneaster salicifolia repandens	compacta	Schinus terebinthifolius
	Hydrangea macrophylla hortensis	Tecoma stans
	"Excelsior"	Weigela "Bristol Snowflake"

PLANTS SHOWING MODERATE TO SEVERE NECROSIS (leaf burn)

Acacia podalyriifolia	Cotoneaster bullata floribunda	Magnolia grandiflora "Majestic Beauty"
Alnus rhombifolia	Dendromecon harfordi rhamnoideis	Philadelphus "Minnesota Snowflake"
Aucuba japonica "Dwarf Female"	Eucalyptus ficifolia	Philadelphus virginialis
Beloperone tomentosa	Gelsemium sempervirens	Potentilla fruticosa "Moonlight"
Berberis "Mentorensis"	Grevillea "Noeli"	Potentilla grandiflora
Ceanothus "Ray Hartman"	Hedera helix "Goldheart"	Quercus agrifolia
Cissus capensis	Leptospermum scoparium "Keatley"	Ternstroemia japonica
Cissus hypoglauca	Liquidambar orientalis	Tristania conferta

IMPORTANT NOTICE — Seller warrants that it is product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.