

Reg # 400-82 **NEW PM-13 LABELING**

174

OMITE®-30W

AGRICULTURAL MITICIDE
RECOMMENDED FOR AGRICULTURAL USE ONLY



COMPOSITION

Active Ingredients: (% by weight)
 Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]* 30.0%
 Inert Ingredients 70.0%
 Total: 100.0%

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749

EPA REG. NO. 400-82

EPA EST. NO.

OMITE is a Registered Trademark of Uniroyal Chemical Company, Inc.

©Copyright 1984 Uniroyal Chemical Company, Inc.

NET WEIGHT

BEST AVAILABLE COPY

SENTENCE
 WHICH WAS
REVISED
UNDERLINED
 ON PAGE 2

NOTIFICATION
 LABEL NOT REVENUED
 DATE JUN 17 1991

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

IF INHALED: Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.

IF ON SKIN: Wash thoroughly with soap and water. Get medical attention.

IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

DANGER

Corrosive, causes eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash hands and face thoroughly with soap and water after use and before eating or smoking. (See use restrictions for protective clothing and clean up requirements)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

REENTRY STATEMENT- EXCEPT FOR HARVESTING

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas until sprays have dried. Protective clothing requirements are outlined below for the indicated reentry.

CROP	EARLY REENTRY WORKER	TIME FOR REENTRY	
		LOW EXPOSURE ACTIVITIES	ALL OTHER ACTIVITIES
Brownberries	when sprays have dried	48 hours	3 days
Grapes	when sprays have dried	48 hours	14 days
Grapes - wine & raisin	when sprays have dried	43 hours	21 days
Grapes - table	when sprays have dried	48 hours	21 days
Citrus	when sprays have dried	48 hours	28 days
Citrus - 10.5 lbs / per acre or less	when sprays have dried	48 hours	28 days
Citrus - at rates more than 10.5 lbs	when sprays have dried	48 hours	28 days
All other crops	when sprays have dried	48 hours	7 days

DIRECTIONS FOR USE

OMITE-30W is a dry, wettable powder for preparing sprays to control brown almond, citrus red, clover, European red, McDaniel, Pacific spider, peach silver, Southern red, strawberry spider, two-spotted spider and Willametta mites.

OMITE-30W is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Always use at least the minimum recommended amount of OMITE-30W per acre. OMITE-30W can be used in concentrate sprays, however, do not reduce the recommended rate per acre.

OMITE-30W performance is best when day temperatures average above 70°F.

1. Early entry is defined as any worker who is directed to enter a reentry restricted area prior to the expiration of any reentry interval that applies to that area. This will include those activities necessitated by unexpected events such as weather, pest conditions, equipment failure, etc. and does not include routine cultural practices.

2. These workers will wear protective clothing to include at least:

- a hat or other suitable head covering
- long-sleeved shirt and long-legged trousers or a coverall type garment (closely woven fabric covering the body including arms and legs).
- chemical resistant gloves.
- shoes and socks.

NOTE: Occupants of enclosed cab vehicles must wear normal work attire.

3. These workers will wear normal work attire to include at least:

- a hat or other suitable head covering.
- long-sleeved shirt and long-legged trousers
- shoes and socks

4. All other activities include those which have high foliar exposure such as pruners, cane turners and tippers (table grapes), etc.

5. See use directions for time of harvest after last application.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to the workers who are expected to be in a treated area or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER, Area to be treated with Propargite on (date of application). Do not enter until sprays have dried. Tractor drivers and irrigators and scouts wearing normal work attire may enter after 48 hours. Do not enter treated areas without protective clothing to perform activities that involve substantial or prolonged exposure with treated foliage for (enter number of days)."

If accidentally exposed, consult the Statement of Practical Treatment portion of this label for first aid recommendations.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

USE RESTRICTIONS

Mixer/loaders must wear goggles or a face shield, chemical-resistant apron, long-sleeved shirt, long pants, and mid-forearm to elbow length chemical-resistant gloves.

Applicators, if applying more than 10 lbs. of OMITE-30W per acre in air blast equipment must be in an enclosed cab. At rates of 10 lbs. or less and if not in an enclosed cab, applicators must wear goggles, wide brimmed hat, a long-sleeved shirt and long pants, and chemical-resistant gloves while applying this pesticide.

Any article of clothing worn while mixing, loading or applying product must be cleaned before re-use. Clothing should be laundered separately from household articles. Clothing that has been drenched or has otherwise absorbed concentrated pesticide must be disposed of in a sanitary landfill, incinerated, or burned if allowed by State and local authorities.

Do not apply this product through any type of irrigation system.

Do not plant any food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite.

Do not feed livestock on cover crops grown among the trees listed on this label.

BEST AVAILABLE COPY

FOR USE ON BEARING CROPS
DOSAGE INSTRUCTIONS (see application notes for specific directions)

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	EARLIEST HARVEST DAYS AFTER APPLICATION
			Ground	Aerial		
ALMONDS (CA & AZ)	clover European red two-spotted spider strawberry spider Pacific spider	5-15 lbs.	50	—	2	28
APPLES ¹	European red two-spotted spider Pacific spider McDaniel	5 - 12 lbs.	500 - 800	—	3	7
APRICOTS (West of Rocky Mts.)	Pacific spider two-spotted spider	5 - 9 lbs.	50	—	2	14
FIGS	two-spotted spider Pacific spider strawberry spider	3 - 6 lbs.	50	—	2	14
GRAPES (West of Rocky Mts.)	Pacific spider strawberry spider Willamette two-spotted spider	5 - 9 lbs.	40	—	2	21
HOPS ¹	two-spotted spider	5 lbs.	100	—	2	14
LEMONS ² (CA & AZ)	citrus red	10 lbs.	100	—	2	21
		11 - 15 lbs.	100	—	2	28
NECTARINES (West of Rocky Mts.)	Pacific spider brown almond	5 - 9 lbs. **	50	20	2	14
ORNAMENTALS CARNATIONS CHRYSANTHEMUMS ³	two-spotted spider	1 - 1 1/2 lbs. per 100 gals. water	Spray to wet	—	—	—
CHRISTMAS TREES & CONIFERS ⁴ (for plantation & nursery use only)	Spider mites	4 1/2 - 7 1/2 lbs (west)	100	10	3	—
		3 - 7 1/2 lbs (east)				
ROSES ³ (field grown)	two-spotted spider Pacific spider Southern red	2 - 5 lbs.	50	—	—	—
OTHER ORNAMENTALS (field & nursery)	two-spotted spider Southern red	1 lbs. per 100 gallons	Spray to wet	—	—	—
PEACHES	European red Pacific spider two-spotted spider strawberry spider McDaniel peach silver	5 - 15 lbs. **	50	20	2	14
PEANUTS	two-spotted spider	3 - 5 lbs.	20	—	2	14
PLUMS PRUNES	European red two-spotted spider Pacific spider	5 - 15 lbs **	50	20	2	12
STRAWBERRIES ⁴ (West of Rocky Mts.)	Pacific spider strawberry spider two spotted spider	5 - 10 lbs.	100	—	3 per 40 day harvest season	3
STRAWBERRIES ⁴ (LA & FL)	Pacific spider strawberry spider two-spotted spider	2 1/2 lbs	100	—	3 per 21 day harvest season	3
WALNUTS (CA only)	European red two-spotted spider	7 - 22 lbs	100	10	2	21

** Use a minimum of 7 1/2 pounds of OMITE-30W per acre when applied by air.

(BEST AVAILABLE COPY)

BEST AVAILABLE COPY

474

FOR USE AFTER HARVEST AND ON NON-BEARING TREES ONLY
Non-bearing trees are those that will not bear fruit within one year of Omite Application

DOSEAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT OMITE-30W PER ACRE	MINIMUM GALLONS SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR
NAVEL ORANGES ² (West of Rocky Mts.)	citrus red	10 - 15 lbs.	100	1
GRAPEFRUIT ² (West of Rocky Mts.)	citrus red	10 - 15 lbs.	100	1
CHEERRIES (West of Rocky Mts.)	European red two-spotted spider Pacific spider McDaniel	8 lbs.	50	—
PEARS ³	European red two-spotted spider	5 - 7 1/2 lbs.	50	For use after harvest only.

APPLICATION NOTES:

1. Apples and Hops: Leaf injury and/or fruit spotting may occur if the following conditions are not observed: Do not apply OMITE-30W with Diazinon WP if air temperatures are expected to exceed 95°F the day of spraying apples and hops.

Do not apply in combination with Guthion emulsifiable concentrate. Do not apply OMITE-30W with foliar spray oils because of incompatibility. See Compatibilities or contact your supplier or Uniroyal for further information.

2. Citrus: OMITE-30W may cause some leaf burn on new growth (feather growth) of lemons.

Foliar oil applications must not follow OMITE-30W treatments within 30 days.

OMITE-30W when used alone may follow foliar oils at an interval of greater than 40 days after an oil application.

Do not use on navel oranges or grapefruit after new fruit is exposed in the spring as fruit spotting may occur.

3. Ornamentals: 1 lb. of OMITE-30W per 100 gallons of water equals 1 ounce (28.4 grams) per 6.25 gallons. Because cultivars and cultural practices differ widely, the user should initially treat only a small portion of his crop to confirm plant safety under his conditions. See list of ornamentals on this label which may show phytotoxicity. Repeat

applications of OMITE-30W are normally necessary for extended control.

4. Strawberries: Do not combine with other chemicals or spray adjuvants, when using on strawberries, as leaf and fruit injury may occur.

California: Do not apply at day temperatures above 80°F as fruit burn may occur.

5. Christmas Trees and Conifers: Higher or lower gallonage can be used depending upon tree size, mite population and equipment used to obtain thorough coverage.

6. Do not use on Asian pears (pear apples) as leaf injury may occur.

COMPATIBILITIES — Except as noted OMITE-30W is compatible with many pesticides (See application notes). Check with your Supplier or Uniroyal for details. Observe all cautions and limitations of labeling of all products used in mixtures.

- Do not use additional spray adjuvants
- OMITE-30W is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of OMITE-30W will be reduced.
- Do not apply in combination with foliar spray oils as phytotoxicity may result.

The following ornamentals may show some signs of phytotoxicity.

PLANTS SHOWING SLIGHT NECROSIS (leaf burn)

Abelia grandiflora	Coccoloba laurifolia	Gardenia jasminoides "Vetchi"	Lagerstroemia "Petite Orchid"
Acer palmatum	Cotoneaster horizontalis	Hedera cuneata	Lagerstroemia "Petite Pinkie"
Alectryon subcinerus	Cotoneaster salicifolia repandens	Hedera "Ghost Tree"	Manisotia repens
Ardisia scabrida	Cotoneaster "Skogsholmen"	Hydrangea macrophylla "Hamburg"	Nyssa sylvatica
Aucuba japonica picturata	Cupressus anacardioides	Hydrangea macrophylla hortensis compacta	Pachysandra terminalis
Aucuba japonica variegata	Eucalyptus cinerea	Hydrangea macrophylla hortensis "Excelsior"	Philadelphus "Glacier"
Besleria grandiflora	Eucalyptus globulus compacta	Hydrangea macrophylla "Mamie's Beauty"	Podocarpus gracilior
Berberis thunbergii atropurpurea	Eucalyptus niphophila	Hydrangea macrophylla "Revelation"	Prunus integrifolia
Berberis thunbergii atropurpurea	Eucalyptus rostrata	Hydrangea macrophylla "Trophée"	Rosa Banksea alba
nana "Crimson Pygmy"	Eucalyptus sideroxylon rosea	Hydrangea macrophylla "White"	Schinus terebinthifolius
Bougainvillea rhodamnia	Eucalyptus torquata	Ilex "Nellie Stevens"	Ternstroemia
Buxus microphylla "Richardi"	Eucalyptus viminalis	Loebachia paniculata	Weigela "Bristol Snowflake"
Cassia mandarinum	Gardenia jasminoides "Mystery"		

PLANTS SHOWING MODERATE TO SEVERE NECROSIS (leaf burn)

Acacia podalyriifolia	Cassia cuneata	Grevillea "Noeli"	Philadelphus virginialis
Alnus rhombifolia	Cassia hypoglauca	Hedera helix "Goldheart"	Prunella frutescens "Violetlight"
Aucuba japonica "Dwarf Female"	Cotoneaster bullata floribunda	Leptodermis scoparium "Kestley"	Picea grandiflora
Beloperone tomentosa	Dendromecon harfordii rhomboides	Liquidambar orientalis	Quercus agrifolia
Berberis "Mertonensis"	Eucalyptus ficifolia	Magnolia grandiflora "Majestic Beauty"	Ternstroemia japonica
Ceanothus "Ray Hartman"	Gelsemium sempervirens	Philadelphus "Minnesota Snowflake"	Ternstroemia conferta

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.