

ACCEPTED

MAY 22 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-83

1y3

# OMITE®-57E

## AGRICULTURAL MITICIDE

Recommended for agricultural use only

### Net contents:

### COMPOSITION

Active Ingredients: (% by weight)

Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]\* ..... 57%

Inert Ingredients: ..... 43%

Total: ..... 100%

\*Contains 5 lbs. active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN DANGER • PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Immediately flush eyes with plenty of water. See a physician.

**IF SWALLOWED:** Do not induce vomiting. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

**IF INHALED:** Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.

**IF ON SKIN:** Wash thoroughly with soap and water. Get medical attention.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive, causes eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing.

### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category E in an EPA chemical resistance category selection chart.

**Applicators and Other Handlers Must Wear:** A long-sleeved shirt & long pants; chemical resistant gloves such as barrier laminate, or nitrile rubber, or neoprene rubber, or viton; shoes plus socks; protective eyewear; chemical resistant headgear for overhead exposure; chemical-resistant apron for mixing and loading. Applicators, if applying more than 72 ounces of OMITE-57E per acre in air blast equipment must be in an enclosed cab. The enclosed cabs must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

### PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame

**UNIROYAL  
CHEMICAL**

2 3

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### GENERAL INSTRUCTIONS

OMITE-57E is a liquid emulsifiable concentrate for preparing sprays to control mites.

**OMITE-57E is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.**

Add recommended dosage (from table) of OMITE-57E to the spray tank, agitate and spray thoroughly to cover foliage and fruit for best results.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 7 days.

Exception: After the first 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- chemical resistant gloves such as barrier laminate or nitrile rubber, or neoprene rubber, or viton
- shoes plus socks
- protective eyewear
- chemical resistant headgear for overhead exposure

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### USE RESTRICTIONS

Do not graze or feed livestock on cover crops grown among the trees listed on this label.

Do not plant leafy vegetables in rotation within 2 months after last application of propargite to cotton or corn.

Do not plant any other food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite.

Do not feed animals apple pomace made from treated apple

CROP	MITES CONTROLLED	TIMING OF APPLICATION	AMOUNT OF COMITE II PER ACRE	GALLONS OF SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
				Ground Application	Aerial Application		
BEANS <sup>1)</sup> (green, lima & dry) (West of Rocky Mts.)	two-spotted spider		2¼ to 3¼ pints	minimum of 20	minimum of 5	2	succulent - 7 dry - 28
COTTON <sup>4)</sup> (East of the Rocky Mts.)	two-spotted spider Pacific spider Strawberry spider	Early <sup>2)</sup>	1¼ pints	15 <sup>3)</sup> to 30	Not recommended	3	50
		Mid Season to Layby	1¼ to 2¼ pints	25 <sup>3)</sup> to 40	5 to 15		
		Layby to Boll Opening	2¼ pints	25 <sup>3)</sup> to 50	5 to 15		
FIELD CORN <sup>5)</sup> & POPCORN	two-spotted spider Banks grass		2¼ pints	minimum of 20	minimum of 2	1	30
PEANUTS <sup>7)</sup>	two-spotted spider		2¼ pints	minimum of 20	minimum of 5	1	14
POTATOES <sup>8)</sup> (Pacific Northwest Only)	two-spotted spider		1½ to 2¼ pints	20 to 50	minimum of 10	2	14
SORGHUM <sup>9)</sup> (East of Rocky Mountains)	two-spotted spider Banks grass		1½ to 2¼ pints	minimum of 20	minimum of 5	1	30

### APPLICATION NOTES

#### BEANS

- Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

#### COTTON

- COMITE II may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
- COMITE II may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE II dosage rate per acre.
- Commerically acceptable mite control may be difficult to obtain where synthetic pyrethroids are or have been used.

#### FIELD CORN

- Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. **Ensure applications are made only when corn leaves are dry.** This is especially true for white corn varieties which are the most susceptible to plant injury.
- In Kansas and Colorado use a minimum of 2 gallons spray solution per acre. In Texas and New Mexico and other states, use a minimum of 5 gallons spray solution per acre.

#### PEANUTS

- When temperatures are greater than 90° F with high humidity some leaf phytotoxicity may occur.

#### POTATOES

- The 2¼ pints per acre rate should be used when the crop is under extreme mite pressure.

#### SORGHUM

- Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE II on a large area.

### COMPATIBILITIES

Under certain conditions a non-penetrating surfactant can be added. Rely on your prior use experience or consult your local surfactant supplier for specific recommendations. Do not use penetrating surfactants.

Do not apply in combination with petroleum based foliar spray oils as phytotoxicity may occur.

COMITE II is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of COMITE II will be reduced.

Except as noted COMITE II is compatible with many pesticides. Observe all cautions and limitation of labeling of all products used in mixtures.

Additional information on compatibility is contained in the Uniroyal chemical product label book. Contact your supplier or Uniroyal representative for this information.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.  
**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.  
**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**IMPORTANT NOTICE** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

\* COMITE is a Registered Trademark of Uniroyal Chemical Company, Inc.

© Copyright 1996, Uniroyal Chemical Company, Inc.