

STORE IN COOL DRY PLACE

CAUTION!

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. DO NOT TAKE INTERNALLY. USE WITH ADEQUATE VENTILATION. AVOID CONTACT WITH SKIN OR EYES.

In case of skin contact, wash with plenty of soap and water. In case of contact with eyes, flush with water. If swallowed, induce vomiting. GET IMMEDIATE MEDICAL ATTENTION.

MOR-CRAN*

A SELECTIVE GRANULAR HERBICIDE FOR THE CONTROL OF NUTGRASS, HORSETAIL AND CERTAIN OTHER WEEDS IN CRANBERRIES.

PRE-EMERGENCE WEED KILLER

COMPOSITION

Active Ingredients (Percent by Weight)	
Sodium N-1 Naphthylphthalate*	8.1
Isopropyl N-(3 Chlorophenyl) Carbamate**	5.0
Inert Ingredients	86.9
Total	100.0

*Equivalent to 7.5% N-1 Naphthylphthalamic acid
**Epoxy IPC (U.S. Patent No. 2,693,723)

MOR-CRAN*

MOR-CRAN*

ACCEPTED

DEC 10 1971

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 700 374

FOR USE IN MASSACHUSETTS ONLY

ACCEPTED

AUG 10 1969

NET 50 POUNDS

another product of



LOT NO.

• Trademark of Uniroyal, Inc.
© Copyright 1969 by Uniroyal, Inc.
ALL RIGHTS RESERVED

MOR-CRAN*

GRANULAR HERBICIDE
FOR CRANBERRIES

GRANULAR HERBICIDE
FOR CRANBERRIES

A selective granular herbicide for the control of the following weeds in cranberries:

Nut Grass	Cut Grass
Horse Tail	Summer Grass
Warty Panicgrass	Mud Rush
Corn Grass	Canada Rush

GENERAL INSTRUCTIONS

1. Preparation of Soil

Area to be treated must be free of all weeds at the time of application. MOR-CRAN usually will not control emerged weeds and grasses.

2. Method of Application

Uniform coverage of soil surface is essential. Apply MOR-CRAN as a dry granular treatment in any suitable air or ground applicator that will insure thorough and uniform coverage.

CRANBERRIES

Apply MOR-CRAN as a broadcast treatment preemergence to the weeds. Make application on well-established plants in the dormant season prior to bud break. From April 1 to May 10 is the optimum time for treatment in Massachusetts.

DIRECTIONS FOR USE

Place MOR-CRAN in the hopper of a granular applicator and apply the required amount of material onto the soil surface before weeds and grasses have emerged. Apply only when plants are dry to avoid contact and sticking with possible injury to fruit buds and loss of yield. Accurate calibration of the applicator is essential for good results.

RECOMMENDED DOSAGE RATE FOR CRANBERRIES

Apply MOR-CRAN at the rate of 100 pounds (13.1 pounds active ingredients) per acre as a broadcast application. Use only on well-established plants in the dormant season prior to bud break.

NOTE 1. In using ground equipment it is suggested that a piece of canvas or burlap be dragged behind the applicator over the plants to brush any granules adhering to the foliage onto the soil surface.

NOTE 2. Good frost protection management should be carried on in the treated areas to minimize the possibility of crop reduction.

NOTE 3. Always apply in the dormant stage before the vines have started the current season's growth.

CAUTION!

1. Container storage and handling: Avoid contamination of foodstuffs, feeds, fertilizers, seeds, plants and pesticides. Do not reuse container. Store in a cool, dry location, avoiding water or oil soaked floors.
2. Do not allow MOR-CRAN to come in contact with such sensitive crops as beets, tomatoes and spinach.
3. Thoroughly brush out equipment after application is completed. If the granular equipment is to be used on sensitive crops it is advisable to run a small amount of some inert material such as sand or cornmeal through the hopper and hoses to remove traces of MOR-CRAN.

IMPORTANT NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

CONTAINER DISPOSAL: BURN OR BURY 18 INCHES AT SAFE DISPOSAL SITE.
UNIROYAL CHEMICAL—Division of UNIROYAL, Inc.—NAUGATUCK, CONNECTICUT 06470

U. S. D. A. Reg. No. 44000

FREIGHT SHIPPING BAG
Meeting requirements of
APPLICABLE FREIGHT CLASSIFICATION
Guaranteed by ST. PETERS PAPER CO.

3-5P-40A

FISH and WILDLIFE
This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on label.