

400-89

2-22-1971

1/2

WARNING! KEEP OUT OF REACH OF CHILDREN.

(For other caution statements see below)

OMITE®-6 E

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the two-spotted spider mite.

Omite-6E is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Mites hit by the spray stop feeding and are killed within 48 to 72 hours, depending on temperature. Omite has some ovicidal activity. The young mites that do hatch from eggs after the spray is applied will be killed by the residue of Omite left on the leaves and fruit.

Add recommended dosage of Omite-6E to the spray tank, add water, agitate and spray thoroughly to cover foliage (both upper and lower leaf surfaces) and fruit for best results. See table for dosage rates.

HOPS -West Coast - For the control of two-spotted spider mites use 20 to 32 fluid ounces per acre.

Add 10 to 16 fluid ounces of Omite-6E in 100 gallons of water and apply at 200 gallons of liquid per acre. Two treatments can be made in one season. Spray can be applied up to 14 days before harvest.

DOSAGE INSTRUCTIONS

CROPS	MITES CONTROLLED	AMOUNT PER ACRE	Amount per 100 Gallons of Water	GALLONS OF SPRAY SOLUTION PER ACRE	TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
			WEST OF ROCKY MOUNTAINS			
Hops	Two-spotted spider	20 fluid ounces - 1 quart	10 - 16 fluid ounces	200	2	14

WARNING

HANDLING: Avoid inhaling vapor or spray mist. Do not get in eyes. May cause eye damage. The wearing of goggles is recommended. In case of contact with eyes, flush with running water for 15 minutes and get medical attention. In case of skin contact, wash thoroughly with warm water and soap. May be harmful if swallowed.

FISH AND WILDLIFE: Keep out of any body of water. This product is toxic to fish.

USE RESTRICTIONS: Do not graze or feed livestock on cover crops grown among the vines listed on this label. Do not feed hop refuse or cones to livestock.

COMPATIBILITY: Omite-6E is compatible with captan, Chlorobenzilate, Cyprax, DDD, Diazinon*, dieldrin, Dikar, Endrin, Ethion, ferbam, fixed neutral copper, Folpet, Gardona, Glyodin, Guthion WP*, Imidan, Karathane, lead arsenate, malathion, maneb, Meta-Systox-R, Methyl parathion, parathion, Phosdrin, Phygon, rotenone, sabadilla, Sevin, sulfur, Systox, TEPP, Thiodan, thiram, Toxaphene, Trithion, zinc oxide, and zineb.

***CAUTION:** Do not use at day temperatures above 95 °F West of the Rocky Mountains.

NOTE: Do not use Omite-6E mixed with alkaline materials such as lime, lime sulfur or Bordeaux mixture. Do not use spring or summer foliar oil sprays before, with, or following Omite-6E. Do not use more than four ounces of additional spray adjuvant per 100 gallons. Do not add any additional spray adjuvants to the spray tank if Omite-6E is mixed with one or more pesticide formulations.

CONTAINER DISPOSAL: Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food.

IMPORTANT NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Net Weight 1 Liquid Gallon

COMPOSITION

Active Ingredient (% by weight)
2-(p-tert-butylphenoxy)cyclohexyl
2-propynyl sulfite*..... 68.1**
Inert Ingredients..... 31.9
TOTAL..... 100.0

**6# Active per gallon

* U.S. Patents 3,272,854; 3,463,859

another product of



2/2

ACCEPTED

FEB 22 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 400-89 SUBJECT
TO ATTACHED COMMENTS.