

400-89

5-7-1973

B7 1/4

RECORDED  
20 JUNE 73

WARNING! KEEP OUT OF REACH OF CHILDREN  
(For other caution statements see back panel)

OMITE (R) -6E  
AN AGRICULTURAL MITICIDE

NET WEIGHT \_\_\_\_\_ LIQUID \_\_\_\_\_

ACCEPTED  
MAY 7 - 1973  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 400-87 SUBJECT  
TO ATTACHED COMMENTS.

COMPOSITION

Active Ingredient (% by weight)	
2-(p-tert-butylphenoxy)cyclohexyl	
2-propynyl sulfite* .....	68.1**
Inert Ingredients .....	31.9
Total .....	100.0

\*\*Contains 6 pounds active Omite per gallon

another product of  
UNIROYAL

\*U.S. Patents 3,272,854; 3,463,859

(R) UNIROYAL'S Registered Trademark for its Miticide

OMITE-6E

AN AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the citrus red mite.

Omite-6E is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. Mites hit by the spray stop feeding and are killed within 48 to 72 hours, depending on temperature. Omite has some ovicidal activity. The young mites that do hatch from eggs after the spray is applied will be killed by the residue of Omite left on the leaves and fruit. Omite performance is best when temperature is 70°F or higher.

Add recommended dosage of Omite-6E to the spray tank, add water, agitate and spray thoroughly to cover foliage (both upper and lower leaf surfaces and fruit for best results).

DIRECTIONS FOR USE

Oranges, lemons - California only

For the control of citrus red mite Panonychus citri (McG)

Use 4 to 6 pints Omite 6E per acre as a dilute spray. Do not use less than 4 pints per acre on mature trees.

Add 6 to 12 fluid ounces Omite 6E to 100 gallons of water and apply 600 to 1,500 gallons of liquid per acre.

Two treatments can be made per season. The last can be applied up to 7 days before harvest.

(Omite-6E may cause some leaf burn on new growth - (feather growth).)

B9 3/4

Omite-6E

AN AGRICULTURAL MITICIDE

NOTE: - Do not use Omite-6E mixed with alkaline materials such as lime, lime sulfur or Bordeaux mixture. Do not use spring or summer foliar oil sprays before, with, or following Omite-6E. Do not use more than four ounces of additional spray adjuvant per 100 gallons. Do not add any additional spray adjuvants to the spray tank if Omite-6E is mixed with one or more pesticide formulations.

HANDLING: - Avoid inhaling spray. Do not get in eyes. May cause eye damage. The wearing of goggles is recommended. In case of contact with eyes, flush with running water for 15 minutes and get medical attention. In case of skin contact, wash thoroughly with warm water and soap. May be harmful if swallowed.

FISH AND WILDLIFE - Keep out of any body of water. This product is toxic to fish.

USE RESTRICTIONS - Do not graze or feed livestock on cover crops.

COMPATIBILITY - Omite-6E is compatible with C-Lite, cygon (Defend), diazinon as wettable powder, dibrom, dieldrin, guthion, karathane, malathion, neutral copper, parathion phosdrin, sabadilla, sevin as wettable powder, sulfur, trithion, urea and zinc oxide.

B10 4/4

OMITE-6E  
AN AGRICULTURAL MITICIDE

CONTAINER DISPOSAL -- Burn or bury container 18 inches below soil surface at save disposal site. Never reuse for food.

IMPORTANT NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

LOT NO.

USDA REGISTRATION NO. 400-

UNIROYAL CHEMICAL, Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770