

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
WARNING**

Causes eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not breathe vapors or spray mist. Do not get in eyes on skin or on clothing. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp and oysters. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

1. **PROHIBITIONS**
Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.
2. **PESTICIDE DISPOSAL**
Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
3. **CONTAINER DISPOSAL**
Triple rinse (or equivalent) and offer for recycling, or disposal in approved landfill or burn, in a safe place.
4. **GENERAL**
Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

OMITE®-6E

AGRICULTURAL MITICIDE

UNIROYAL

ACCEPTED

1982

COMPOSITION

Active Ingredient (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	68.1%**
Inert Ingredients	31.9%
Total	100.0%

*U.S. Patent Nos. 3,272,854; 3,463,859

**6 lbs. Active per Gallon

**KEEP OUT OF REACH OF CHILDREN.
WARNING**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, INC.
Naugatuck, Connecticut 06770
EPA Establishment No.
EPA Registration No. 400-89-AA

NET CONTENTS:

12-29-81

Omite®-6E is a liquid, pourable concentrate for preparing sprays to control European red, Pacific spider, southern red, strawberry spider, and two-spotted spider mites.

Omite-6E is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

Add recommended dosage (from table) of Omite-6E to the spray tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

See back side for dosage instructions chart and application notes.

AL USUARIO:

Si usted no puede leer o entender ingles no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TO THE USER:

If you cannot read or understand English, do not use this product until the label has been fully explained to you.

USE RESTRICTIONS: Do not graze or feed livestock on cover crops grown among the trees and vines listed on this label. Do not feed hop refuse or cones to livestock. However, spent hops from the extract operation may be fed. Do not feed mint spent hay to livestock.

COMPATIBILITIES: Omite-6E is compatible with most pesticides (see caution statements). Check with your supplier for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CAUTION:

- A. Leaf injury may occur if the following conditions are not observed:
 1. Do not apply Omite-6E with Diazinon WP if air temperatures are expected to exceed 95°F the day of spraying hops.
 2. Omite-6E is not compatible with foliar spray oils. (Further information available from supplier.)
 3. Do not apply Omite-6E to cranberries if air temperatures are above 85°F.
- B. Omite-6E is not compatible with alkaline materials such as lime, lime sulfur or Bordeaux. The effectiveness of Omite will be reduced.

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	AMOUNT PER ACRE	Gallons of Spray Solution Per Acre		NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
			Ground Application	Aerial Application		
Almonds (California only)	Clover European red strawberry spider two spotted spider pacific spider	3-4 pints		4-1	2	28
Apricots (California only)	two spotted spider Pacific spider	2-3 pints		minimum of 10 gallons	2	post harvest only
Cranberries ² (Massachusetts only)	southern red mite	30-60 ^{3,4} fluid ounces	100		2	14
Field Corn (California only)	two spotted spider	2 pints	20-50	minimum of 10 gallons	1	56
Hops	two spotted spider	2 pints	200		2	14
Mint	two spotted spider	2-3 pints	20-50	minimum of 10 gallons	2	14
Potatoes (Pacific Northwest only)	two-spotted spider	2-3 pints	20-50	minimum of 10 gallons	2	14
Walnuts	two spotted spider European red	2-9 pints	500-1500	minimum of 20 gallons	2	14

APPLICATION NOTES: — ALMONDS

- One application of Omite 6E may be applied by air to almonds over one application of Omite-30W.

APPLICATION NOTES: — CRANBERRIES

- Omite-6E may be applied through a sprinkler irrigation system to cranberries.
- 30-60 ounces Omite-6E may be applied to cranberries once prior to bloom followed by a second application of 30 ounces any time prior to 14 days harvest.
- Regardless of the size of the irrigation system, if Omite-6E is used at the recommended rate per acre, the ratio of diluent to product in the slurry tank should be approximately 4:1.

APPLICATION NOTES: — WALNUTS

- Use 3 to 4 pints of Omite-6E in a minimum of 20 gallons of water per acre when applied by air.

IRRIGATION USE PRECAUTIONS

- Apply the product only through systems containing anti-siphon and check valves which will prevent water source contamination and overflow of the slurry tank, and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.
- Constant agitation must be maintained in the chemical supply tank during the entire period of insecticide application.
- Inject the product with a positive displacement pump into the main line ahead of a right angle turn, to insure adequate mixing.
- Application of more than label recommended quantities of irrigation water per acre may result in decreased product performance by removing the chemical from the zone of effectiveness.
- Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.
- Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of more dilute slurry per hour.
- Where sprinkler distribution patterns do not overlap sufficiently unacceptable mite control may result. Where sprinkler distribution patterns overlap excessively crop injury may result.

IMPORTANT NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.