

PA21 400-92

**STATEMENTS
HAZARDS TO HUMANS
DANGER**

Corrosive, causes eye damage and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid inhaling of spray mist or dust.
Do not use or store near food or feed.

During commercial exposure in seed treatment and loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety and Health Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under provisions of 30 CFR, Part 11. Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work; wash all clothing with soap and water before reusing. Consumption of alcoholic beverages increases the toxic effects of thiram.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a dry location
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

CONTAINER DISPOSAL:

Bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke

Fiber Drum: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contami-

VITAVAX-T[®]


FUNGICIDE
(VITAVAX WITH THIRAM)



COMPOSITION

| | |
|--|---------------|
| Active Ingredients (% by weight) | |
| Carbendazim (5-benzimidazol-2-ylmethyl-1,4-oxathiazol-2-ylmethyl-1H-imidazole) | 37.5% |
| Thiram (tetramethylthiuram disulfide) | 37.5% |
| Inert Ingredients | 25.0% |
| Total | 100.0% |

*U.S. Patents 3,249,499 - 3,393,202 - 3,454,391



KEEP OUT OF REACH OF CHILDREN. DANGER


STATEMENT OF PRACTICAL TREATMENT

- If swallowed: Induce vomiting and see a physician
 - If inhaled: See a physician if an abnormal reaction occurs
 - If on skin: Wash thoroughly with soap and water.
 - If in eyes: Immediately flush eyes with plenty of water and get medical treatment
- See left side panel for additional precautionary statement

CHEMICAL GROUP — Uniroyal, Inc.
Middlebury, CT 06749
EPA Establishment No.
EPA Registration No. 400-92

VITAVAX is a Registered Trademark of UNIROYAL, Inc.
Copyright 1985, Uniroyal, Inc.

NET WEIGHT.



DIRECTIONS FOR USE

VITAVAX-T Fungicide is a wettable powder seed treatment for use as a dry dust, spray mix, or slurry seed treatment. In liquid systems, a thin paste should first be made using a portion of the required water and the recommended amount of VITAVAX-T Fungicide. Stir this paste into the remainder of the water and provide enough agitation to maintain a uniform water suspension.

BARLEY, CORN, OATS, AND WHEAT

VITAVAX-T Fungicide combines the systemic activity of carboxin with the surface action of thiram to control various seed and seedling diseases of barley, corn, oats and wheat. It is particularly effective against *Rhizoctonia solani*.

In addition, crown smut, covered smut, flag smut, common bunt and barley stripe are controlled by VITAVAX-T Fungicide.

DOSAGE INSTRUCTIONS

| Crop | Diseases Controlled | Amount of VITAVAX-T Fungicide per 100 Pounds of Seed |
|--------|--|--|
| Barley | seed rot seedling blight Barley Stripe crown smut covered smut | 3.4 ounces |
| Corn | seed rot seedling blight | 4 ounces |
| Oats | seed rot seedling blight crown smut | 3.4 ounces |
| Wheat | seed rot seedling blight crown smut flag smut | 3.4 ounces |

USE RESTRICTIONS — Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

HANDLING — Care must be exercised in handling of treated seed.

SEED LABELING — This seed is treated with VITAVAX-T Fungicide. Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas for six weeks after planting. Add appropriate information for any other chemical treatments.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instruction, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.