

137  
ACCEPTED

CAUTION! KEEP OUT OF REACH OF CHILDREN.  
(See back panel for additional cautions)

400-101

TEST DOCUMENT AVAILABLE

ROYALTAC\*  
CONTACT TOBACCO SUCKER CONTROL AGENT

NET WT. 1 LIQUID GALLON

COMPOSITION

ACTIVE INGREDIENTS (by weight)\*\*

n-Decanol .....	78.4%
n-Octanol .....	0.4%
INERT INGREDIENTS .....	21.2%
TOTAL .....	100.0%

SHAKE WELL BEFORE USING

another product of

\*\*Weight of Active Ingredients  
is 5.70 pounds per gallon.

UNIROYAL

\*Trademark of UNIROYAL, Inc.  
EPA Registration No. 400-101

2

GENERAL INFORMATION

ROYALTAC is a contact sucker control agent. Direct contact of ROYALTAC with the suckers is essential. The spray should wet sides of the stalk of the plant and contact all small immature suckers to control them.

ROYALTAC may be used for the control of sucker growth on all types of tobacco.

CAUTION

1. Avoid inhalation of spray mist. In case of eye contact, flush immediately with water.
2. Application rates of ROYALTAC concentrations higher than recommended levels may cause leaf damage.
3. Shake well before using. After adding ROYALTAC to the sprayer, start up and maintain good agitation of the diluted solution.
4. Do not mix ROYALTAC with other chemicals.
5. Container Disposal - Chop holes top and bottom. Bury 18 inches below soil surface at safe disposal site. Never reuse for food or water.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

3  
29

#400-101

DIRECTION FOR USE

TIME OF APPLICATION

ROYALTAC can be applied either before or after topping, preferably just after buttons have elongated and before flowers start to open. At this stage very few suckers are large enough to require removing by hand. Best results are usually obtained by spraying with ROYALTAC and then topping within a week.

If tobacco is topped first, apply ROYALTAC within two days after topping. Any suckers larger than one inch should be removed while topping.

Do not spray during the middle of the day in very hot weather or leaf burn may occur.

Injury may also result if ROYALTAC is applied to the underside of the tobacco leaf; thus, application in winds high enough to turn over the top leaves is not recommended.

RATE OF APPLICATION

For use on FLUECURED TOBACCO

Mix 1½ gallons of ROYALTAC in 50 gallons of water. Apply 40-60 gallons of diluted product per acre. Then follow in 7 to 10 days by Royal MH-30 or MH-30 systemic sucker control agents at dosages recommended on the label.

For use on BURLEY, MARYLAND and CIGAR TOBACCO.

Mix 2 gallons of ROYALTAC in 50 gallons of water. Apply 40-60 gallons of diluted product per acre.

28