

11/10/11

For DARK TOBACCO, use 7-8 ounces of ROYALTAC per gallon of water. If applied by hand, one gallon of diluted solution will treat 130 to 190 plants.

If a power sprayer is used, 40 to 60 gallons of diluted product should be applied per acre of tobacco.

METHOD OF APPLICATION

When applied with power equipment, three nozzles per row should be used. One nozzle should be directed downward over the center of the row and one should be 9 inches on either side directed at or slightly above the top of the stalk. The diluted ROYALTAC should be applied to the topped tobacco as a coarse spray from a height of 12 to 16 inches above the top of the stalk. It is recommended that boom pressure be kept at 20-25 pounds and should never exceed 30 pounds.

When a power sprayer is used be sure the solution is thoroughly mixed before the spraying operation begins; maintain good agitation.

If you use a hand held or back pack sprayer, the diluted emulsion should be applied at a rate of 2/3 to one ounce per plant (or enough to insure run down to the bottom of the plant). A coarse spray is recommended, directed downward at the top of the stalk from 6 to 8 inches above the top leaves. Very little tank pressure is required, and in no case should more than about 20 pounds be used.

NUMBER OF APPLICATIONS

This application should be followed in 7 to 10 days by ROYAL MH-30 or MH-30 systemic sucker control agents at the dosage recommended on the label. A second ROYALTAC application may be made 10-14 days after the first ROYALTAC application if a systemic material is not used (except on flue cured tobacco).

On fluecured tobacco apply ROYALTAC then MH-30 or ROYAL MH-30 7 to 10 days later.

LOT NO. _____

1972 UNIROYAL, Inc. All Rights Reserved
Printed in U.S.A.

372-1G

UNIROYAL CHEMICAL - Division of UNIROYAL, Inc. Naugatuck Connecticut 06770