

DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not inhale dust or spray. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements see back panel.

ACCEPTED
400-104
JAN 3 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. _____ SUBJECT
TO ATTACHED COMMENTS.

COMITE™

AGRICULTURAL MITICIDE
(EAST OF THE ROCKY MOUNTAINS)

COMITE™

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus and two-spotted spider mites.

Comite is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of Comite to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

Crops	Mites Controlled	Amount of Comite Per 100 Gallons of Water	Gallons Spray Solution Per Acre ²	Total No. of Sprays Per Year	Days Before Harvest (Last Spray)
Oranges and Grapefruit	citrus red Texas citrus six-spotted citrus rust false spider	5-gal. fluid ounces	600-1500 ¹	2	7

¹ Amount of COMITE per 500 Gallon Tank — 25-40 Fluid Ounces
² Total Amount of COMITE Per Acre — 30-120 Fluid Ounces
³ Comite may be applied as a complete coverage spray in Texas at a rate of 1½ pints/250 gallons of spray solution per acre.

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total No. of Sprays Per Year	Last Spray
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches ⁴	1 pint	15 ⁵ to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1½ pints	25 ⁵ to 40 gallons	5 to 15 gallons		
		Over 24 inches	1½ to 2 pints	25 ⁵ to 50 gallons	5 to 15 gallons		

APPLICATION NOTES
4. COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.
5. COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

WARNING!

This product is toxic to fish and shrimp. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Use Restrictions — On cotton, apply only before bolls open. Do not feed treated foliage or cotton trash to livestock. Do not graze or feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES

COMITE is compatible with most pesticides (see caution statements). Check with your supplier for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CAUTION!

A. On citrus, leaf injury may occur if Comite is tank mixed with oil or applied within two weeks prior to or following an oil application.
B. Do not mix Comite with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

CONTAINER DISPOSAL — Bury container 18 inches below soil surface at safe disposal site. Never reuse.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ONE LIQUID GALLON

COMPOSITION

Active Ingredient (% by weight) Propargite	
2-(p-tert-butylphenoxy)cyclohexyl	
2-propynyl sulfite*	75.0%**
Inert Ingredients	25.0%
Total	100.0%

*U. S. Patents 3,272,854; 3,463,859

**Contains 6.75 pounds COMITE per gallon.



EPA Reg. No. 400-104-AA

EPA Est. No.

UNIROYAL CHEMICAL, Division of UNIROYAL, Inc., Naugatuck, Conn. 06770

©1974 Uniroyal, Inc. All Rights Reserved.

Lot No. _____

11-5-74-1G

400-104

1-3-1975

11