

DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage.
Do not get in eyes, on skin or on clothing.
Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not inhale dust or spray. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements see back panel.

COMITE™

AGRICULTURAL MITICIDE
(EAST OF THE ROCKY MOUNTAINS)

ACCEPTED
NOV 03 1975
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended, and the provisions
thereof, this product is hereby
registered.
EPA Reg. No. 400-104-AA

ONE LIQUID GALLON

COMPOSITION

Active Ingredient, by weight	Propargite
2-propargite butylphenoxycyclohexyl	75.0 %**
Inert Ingredients	25.0 %
Total	100.0

U.S. Patents 3,272,894; 3,463,659

Manufactured by COMITE per gallon

UNIROYAL

EPA Reg. No. 400-104-AA

EPA Est.

COMITE™

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus, and two-spotted spider mites.

Comite is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table of Comite to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

Crop	Mites Controlled	Amount of Comite per Acre	Timing of Application	Days Before Harvest
Oranges and Grapefruit	Citrus red, Texas citrus, false spider, Pacific spider, six-spotted, strawberry spider	4.0 lb. per acre	Full bloom	14

APPLICATION NOTES — Oranges and Grapefruit

- Amount of COMITE per 500 Gallon Tank = 25.40 Fluid ounces.
- Total Amount of COMITE Per Acre = 30.120 Fluid ounces.
- Comite may be used in a tank with other pesticides at a rate of 10 parts COMITE to 90 parts of other pesticides.

Crop	Mites Controlled	Amount of Comite per Acre	Timing of Application	Days Before Harvest
Field corn	Two-spotted spider, Banks grass	4.0 lb. per acre	Full bloom	14

APPLICATION NOTES — Corn

- COMITE should be applied by ground application equipment, or by aircraft, at a rate of 10 parts COMITE to 90 parts of other pesticides.
- COMITE may be used in a tank with other pesticides at a rate of 10 parts COMITE to 90 parts of other pesticides.
- COMITE may be used in a tank with other pesticides at a rate of 10 parts COMITE to 90 parts of other pesticides.

Crop	Mites Controlled	Amount of Comite per Acre	Timing of Application	Days Before Harvest
Field corn	Two-spotted spider, Banks grass	4.0 lb. per acre	Full bloom	14

APPLICATION NOTES — Corn

- COMITE should be applied by ground application equipment, or by aircraft, at a rate of 10 parts COMITE to 90 parts of other pesticides.
- COMITE may be used in a tank with other pesticides at a rate of 10 parts COMITE to 90 parts of other pesticides.
- COMITE may be used in a tank with other pesticides at a rate of 10 parts COMITE to 90 parts of other pesticides.

FISH AND WILDLIFE — This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Use Restrictions — On cotton, apply only before bolls open. Do not feed treated forage to livestock. Do not graze livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES

COMITE is compatible with most pesticides. See caution statements. Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CAUTION!

- On citrus, leaf injury may occur if Comite is tank mixed with oil or applied within two weeks prior to or following an oil application.
- Do not mix Comite with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum-type solvents.

CONTAINER DISPOSAL — Destroy empty container and bury at an approved disposal site.

UNIROYAL CHEMICAL Division of UNIROYAL, Inc., Naugatuck, Conn. 06770

© 1974 Uniroyal, Inc. All Rights Reserved

Lot No.