



DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not inhale dust or spray. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements see back panel.

COMITE™

AGRICULTURAL MITICIDE
(EAST OF THE ROCKY MOUNTAINS)

ACCEPTED
NOV 03 1975
Under the Federal Insecticide, Fungicide, and Plant Disease Act,
as amended by the FIFRA Act of 1972
EPA Reg. No. 400-104

ONE LIQUID GALLON

COMPOSITION

Active Ingredient (% by weight) Propargite	
2-(p-tert-butylphenoxy)cyclohexyl	
2-propynyl sulfite*	75.0%**
Inert Ingredients	25.0%
Total	100.0%

*U.S. Patents 3,272,854; 3,463,859

**Contains 6.75 pounds COMITE per gallon.



EPA Reg. No. 400-104-AA
EPA Est.

COMITE™

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus and two-spotted spider mites.

Comite is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of Comite to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

Crops	Mites Controlled	Amount of Comite Per 100 Gallons of Water	Gallons Spray Solution Per Acre ¹	Total No. of Sprays Per Year	Days Before Harvest (Last Spray)
Oranges and Grapefruit	citrus red Texas citrus six-spotted citrus rust false spider	5-8 ¹ fluid ounces	600-1500 ³	2	7

APPLICATION NOTES — Oranges and Grapefruit
1. Amount of COMITE per 500 Gallon Tank — 25-40 Fluid Ounces
2. Total Amount of COMITE Per Acre — 30-120 Fluid Ounces
3. Comite may be applied as a complete coverage spray in Texas at a rate of 1 1/2 pints/250 gallons of spray solution per acre.

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total No. of Sprays Per Year	Last Spray
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches ⁴	1 pint	15 ⁵ to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1 1/2 pints	25 ⁵ to 40 gallons	5 to 15 gallons		
		Over 24 inches	1 1/2 to 2 pints	25 ⁵ to 50 gallons	5 to 15 gallons		

APPLICATION NOTES — Cotton
4. COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.
5. COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

Crop	Mites Controlled	Amount of Per Acre	Total No. of Sprays Per Year	Days Before Harvest (Last Spray)
Field corn	two-spotted spider Banks grass	2 ⁶ pints	1	30

APPLICATION NOTES — Corn
6. COMITE should be applied by ground application equipment in a minimum of 20 gallons of spray solution per acre or by air in a minimum of 2 gallons of spray solution per acre.

FISH AND WILDLIFE — This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

Use Restrictions — On cotton, apply only before bolls open. Do not feed treated foliage or cotton trash to livestock. Do not graze or feed livestock on cover crops grown among the trees listed on this label.

COMPATIBILITIES
COMITE is compatible with most pesticides (see caution statements). Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CAUTION!

- A. On citrus, leaf injury may occur if Comite is tank mixed with oil or applied within two weeks prior to or following an oil application.
- B. Do not mix Comite with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

CONTAINER DISPOSAL — Destroy empty container and bury at an approved disposal site.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

UNIROYAL CHEMICAL, Division of UNIROYAL, Inc., Naugatuck, Conn. 06770

©1974 Uniroyal, Inc. All Rights Reserved.

Lot No. _____

8-25-75-1G

400-104

11-30-1975

1/2

DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not inhale dust or spray. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements see back panel.

COMITE™

AGRICULTURAL MITICIDE
West of the Rocky Mountains

ACCEPTED

NOV 03 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for use as pesticide registered under EPA Reg. No. 400-104

UNIROYAL

EPA Reg. No. 400-104-AA

EPA Est.

5 LIQUID GALLONS

COMPOSITION

Active Ingredient (% by weight) Propargite	
2-(p-tert-butylphenoxy)cyclohexyl	
2-propynyl sulfite*	75.0%**
Inert Ingredients	25.0%
Total	100.0%

*U.S. Patents 3,272,854; 3,463,859

**Contains 6.75 pounds COMITE per gallon.

COMITE™

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, two-spotted, Pacific and strawberry spider mites. Comite is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces is necessary for effective control. Add recommended dosage (from table) of Comite to the spray tank, agitate and spray thoroughly to cover foliage for best results.

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Solution Per Acre		Total Number of Sprays Per Year	Last Spray
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches ⁽¹⁾	1 pint	15 ⁽²⁾ to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1½ pints	25 ⁽²⁾ to 40 gallons	5 to 15 gallons		
		Over 24 inches	1¾ to 2 pints	25 ⁽²⁾ to 50 gallons	5 to 15 gallons		

APPLICATION NOTES — COTTON

- COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.
- COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

Crop	Mites Controlled	Amount Per Acre	Gallons Spray Solution Per Acre	Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
Potatoes (Pacific Northwest Only)	two-spotted spider	1½ - 2½ ⁽¹⁾ pints	20 - 50 ⁽⁴⁾	2	14

APPLICATION NOTES — POTATOES

- The 2½ pints per acre rate should be used when the crop is under extreme mite pressure.
- COMITE can be applied by air to potatoes in a minimum of 10 gallons of spray solution per acre.

Crop	Mites Controlled	Amount Per Acre	Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
Field corn	two-spotted spider Banks grass	2 ⁽³⁾ pints	1	30

APPLICATION NOTES — CORN

- COMITE should be applied by ground application equipment in a minimum of 20 gallons of spray solution per acre or by air in a minimum of 2 gallons of spray solution per acre.

FISH AND WILDLIFE — This product is toxic to fish and shrimp. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

USE RESTRICTIONS — Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.

COMPATIBILITIES

Comite is compatible with most pesticides (see caution statements). Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

CAUTION!

Do not mix Comite with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

SPRAY ADJUVANT — FOR USE ON COTTON OR CORN ONLY

UNIROYAL SPRINT-38 is recommended for best results with COMITE. The following rates are suggested:

Ground Sprays	- 1 pt. SPRINT-38 per 100 gallons water
Aerial Sprays	- 2 pts. SPRINT-38 per 100 gallons water
Concentrate Sprays	- 2 pts. SPRINT-38 per 100 gallons water

Spray adjuvants other than SPRINT-38 should not be used.

CONTAINER DISPOSAL — Destroy empty container and bury at an approved disposal site.

IMPORTANT NOTICE - Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

2/8