

## **PLEASE NOTE**

**This image contains more than one label approved for this product on this date.**

**DANGER! KEEP OUT OF REACH OF CHILDREN.**

Corrosive. Causes skin and eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not inhale dust or spray. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements see back panel.

# COMITE®

AGRICULTURAL MITICIDE  
(EAST OF THE ROCKY MOUNTAINS)

**ACCEPTED**  
JUL 7 1976  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-104-104



EPA Reg. No. 400-104-AA  
EPA Est. No.

**NET CONTENTS:**

COMPOSITION

Active Ingredient (% by weight) Propargite  
2-(p-tert-butylphenoxy)cyclohexyl  
2-propynyl sulfite\* ..... 75.0%\*\*  
Inert Ingredients ..... 25.0%  
Total ..... 100.0%

\*U.S. Patents 3,272,854; 3,463,859

\*\*Contains 6.75 pounds COMITE per gallon.

# COMITE®

AGRICULTURAL MITICIDE

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus and two-spotted spider mites.

Comite is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of Comite to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

**DOSAGE INSTRUCTIONS**

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches (1)	1 pint	15 (2) to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1 1/2 pints	25 (2) to 40 gallons	5 to 15 gallons		
		Over 24 inches	1 1/2 to 2 pints	25 (2) to 50 gallons	5 to 15 gallons		
Field corn	two-spotted spider Banks grass	X	2 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
Grain Sorghum (3)	two spotted spider Banks grass	X	1 1/2 to 2 pints	minimum of 20 gallons	minimum of 5 gallons	1	30 (3)
Oranges and Grapefruit	citrus red Texas citrus six-spotted citrus rust false spider	X	30-120 Fluid Ounces (4)	600-1500 gallons	X	2	7

**APPLICATION NOTES — Cotton**

1. COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.
2. COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

**APPLICATION NOTES — Grain Sorghum**

4. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check for phytotoxicity before using COMITE on a large area.
5. SILAGE STAGE. Grain should not be harvested within 60 days of spraying COMITE. Do not plant rotational crops within 6 months of the last treatment on grain sorghum.

**APPLICATION NOTES — Oranges and Grapefruit**

6. Amount of COMITE per 500 Gallon Tank — 25-40 Fluid Ounces.

**FISH AND WILDLIFE** — This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

**Use Restrictions** — On cotton, apply only before bolls open. Do not feed treated foliage or cotton trash to livestock. Do not graze or feed livestock on cover crops grown among the trees listed on this label.

**COMPATIBILITIES**

COMITE is compatible with most pesticides (see caution statements). Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

**CAUTION!**

- A. On citrus, leaf injury may occur if Comite is tank mixed with oil or applied within two weeks prior to or following an oil application.
- B. Do not mix Comite with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

**CONTAINER DISPOSAL** — Destroy empty container and bury at an approved disposal site.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

UNIROYAL CHEMICAL, Division of UNIROYAL, Inc., Naugatuck, Conn. 06770

400-104

7-7-1976

11

**NEXT**

**LABEL**

**DANGER! KEEP OUT OF REACH OF CHILDREN.**

Corrosive. Causes skin and eye damage.  
Do not get in eyes, on skin or on clothing.  
Wear goggles or face shield and rubber gloves when handling.  
Harmful if swallowed. Do not inhale dust or spray.  
Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements, see right panel.

**ACCEPTED**  
**JUL 7 1976**  
Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the purpose of registration under EPA Reg. No. 400-104

**COMITE®**  
AGRICULTURAL MITICIDE  
(West of the Rocky Mountains)

**Net Contents:**

**COMPOSITION**

Active Ingredient: (% by weight) Propargite	
2-(p-tert-butylphenoxy)cyclohexyl	
2-propynyl sulfite*	75% **
Inert Ingredients:	25%
Total:	100%

\*U.S. Patents 3,272,854 - 3,463,859

\*\*Contains 6.75 pounds COMITE per gallon.



EPA Reg. No. 400-104-AA

EPA Est.

©1974 UNIROYAL, Inc. All Rights Reserved.

6/76/5

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, two-spotted, Pacific and strawberry spider mites.

COMITE is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces is necessary for effective control.

sary for effective control.

Add recommended dosage (from table) of COMITE to the spray tank, agitate and spray thoroughly to cover foliage for best results.

**DOSAGE INSTRUCTIONS**

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
				Ground Application	Aerial Application		
Beans (green and snap lima and dry)	two-spotted spider	X	2-3 pints	minimum of 20 gallons	minimum of 5 gallons	2	7 <sup>(1)</sup>
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches <sup>(2)</sup>	1 pint	15 <sup>(2)</sup> to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1 1/2 pints	25 <sup>(2)</sup> to 40 gallons	5 to 15 gallons		
		Over 24 inches	1 1/2 to 2 pints	25 <sup>(2)</sup> to 50 gallons	5 to 15 gallons		
Field corn	two-spotted spider Banks grass	X	2 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
Potatoes (Pacific Northwest Only)	two-spotted spider	X	1 1/2 to 2 1/2 <sup>(1)</sup> pints	20 to 50 gallons	minimum of 10 gallons	2	14

**APPLICATION NOTES — BEANS**

- Succulent beans. Days before harvest (last spray) for dry beans is 28 days. Do not plant rotational crops within 6 months of the last treatment on beans.

**APPLICATION NOTES — COTTON**

- COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.

- COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

**APPLICATION NOTES — POTATOES**

- The 2 1/2 pints per acre rate should be used when the crop is under extreme mite pressure.

**FISH AND WILDLIFE** — This product is toxic to fish and shrimp. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

**USE RESTRICTIONS —**

- Cotton — Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.
- Beans — Do not feed or forage treated bean vines or bean trash after harvesting.

**COMPATIBILITIES** — COMITE is compatible with most pesticides (see caution statements). Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

**CAUTION:**

Do not mix COMITE with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

**SPRAY ADJUVANT — FOR USE ON COTTON OR CORN ONLY**

- Ground Sprays — 1 pt. SPRINT-38 per 100 gallons water
- Aerial Sprays — 2 pts. SPRINT-38 per 100 gallons water
- Concentrate Sprays — 2 pts. SPRINT-38 per 100 gallons water

Spray adjuvants other than SPRINT-38 should not be used.

**CONTAINER DISPOSAL** — Destroy empty container and bury at an approved disposal site.

**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

401-004

9261-6-7

1/1