

DANGER! KEEP OUT OF REACH OF CHILDREN.

Corrosive. Causes skin and eye damage.
Do not get in eyes, on skin or on clothing.
Wear goggles or face shield and rubber gloves when handling.
Harmful if swallowed. Do not inhale dust or spray.
Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

For other precautionary statements, see right panel.

ACCEPTED

FEB 25 1977

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 400-104

COMITE®
AGRICULTURAL MITICIDE
(West of the Rocky Mountains)

Insert →

2. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.


Net Contents:

COMPOSITION

Active Ingredient: (% by weight) Propargite 2-(p-tert-butylphenoxy)cyclohexyl 2-propynyl sulfite*	75%**
Inert Ingredients:	25%
Total:	100%

*U.S. Patents 3,272,854 - 3,463,853

**Contains 6.75 pounds COMITE per gallon.



EPA Reg. No. 400-104-AA

EPA Est.

© 1974 UNIROYAL, Inc. All Rights Reserved.

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, two-spotted, Pacific and strawberry spider mites.

COMITE is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces is necessary for effective control.

sary for effective control.

Add recommended dosage (from table) of COMITE to the spray tank, agitate and spray thoroughly to cover foliage for best results.

DOSAGE INSTRUCTIONS

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
				Ground Application	Aerial Application		
Beans (green time and dry)	two-spotted spider	X	2-3 pints	minimum of 20 gallons	minimum of 5 gallons	2	7 ¹¹
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches ⁽²⁾	1 pint	15 ⁽²⁾ to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1 1/2 pints	25 ⁽²⁾ to 40 gallons	5 to 15 gallons		
		Over 24 inches	1 1/2 to 2 pints	25 ⁽²⁾ to 50 gallons	5 to 15 gallons		
Field corn	two-spotted spider Banks grass	X	2 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
Potatoes (Pacific Northwest Only)	two-spotted spider	X	1 1/2 to 2 1/2 ⁽¹⁾ pints	20 to 50 gallons	minimum of 10 gallons	2	14

APPLICATION NOTES — BEANS

1. Succulent beans. Days before harvest (last spray) for dry beans is 28 days. Do not plant rotational crops within 6 months of the last treatment on beans.

tative before spraying cotton less than 10 inches in height.

4. COMITE may be used in concentrate sprayers. Use only sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

APPLICATION NOTES — COTTON

3. COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.

APPLICATION NOTES — POTATOES

5. The 2 1/2 pints per acre rate should be used when the crop is under extreme mite pressure.

FISH AND WILDLIFE — This product is toxic to fish and shrimp. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

SPRAY ADJUVANT — FOR USE ON COTTON

Ground Sprays —
1 pt. SPRINT-38 per 100 gallons water
Aerial Sprays —
2 pts. SPRINT-38 per 100 gallons water
Concentrate Sprays —
2 pts. SPRINT-38 per 100 gallons water

USE RESTRICTIONS —

Cotton — Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.
Beans — Do not feed or forage treated bean vines or bean trash after harvesting.

Spray adjuvants other than SPRINT-38 should not be used.

CONTAINER DISPOSAL — Destroy empty container and bury at an approved disposal site.

COMPATIBILITIES — COMITE is compatible with most pesticides (see caution statements). Check with your supplier or Uniroyal for details. Observe all cautions and limitations on labeling of all products used in mixtures.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION!

Do not mix COMITE with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

400-104
2-25-1977