

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
DANGER**

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used, or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

3. CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling, or disposal in approved landfill or bury in a safe place.

4. GENERAL

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

COMITE®

AGRICULTURAL MITICIDE



COMPOSITION

Active Ingredient: (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	73.0%
Inert Ingredients:	27.0%
Total:	100.0%

*U.S. Patents 3,272,854 - 3,463,859
**Contains 6.55 pounds COMITE per gallon.

ACCEPTED
NOV 09 1977
Under the Federal Insecticide, Fungicide, and Plant Disease Act, as amended, this product is registered under EPA Reg. No. 400-104

**KEEP OUT OF REACH OF CHILDREN.
DANGER**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.
Naugatuck, Connecticut 06770
EPA Establishment No.
EPA Registration No. 400-104-AA

NET WEIGHT

9-9-77

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus, and two-spotted spider mites.

COMITE is not systemic in action; therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of COMITE to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

See back side for dosage instructions chart and application notes.

USE RESTRICTIONS —

Cotton — Apply only before bolls open. Do not feed treated foliage of cotton trash to livestock.

Beans — Do not feed or forage treated bean vines or bean trash after harvesting.

Do not graze or feed livestock on cover crops grown among the trees listed on this label.

General — Do not mix COMITE with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

©1974 UNIROYAL, Inc. All Rights Reserved.

400-104

11-9-1977

1/2

DOSAGE INSTRUCTIONS

Crop	Mites Controlled	Height of Cotton	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	10 to 18 inches ⁽¹⁾	1 pint	15 ⁽²⁾ to 30 gallons	Not recommended	3	Before bolls open
		14 to 24 inches	1½ pints	25 ⁽²⁾ to 40 gallons	5 to 15 gallons		
		Over 24 inches	1½ to 2 pints	25 ⁽²⁾ to 50 gallons	5 to 15 gallons		
Field corn ⁽³⁾	two-spotted spider Banks grass	X	2 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
Grain Sorghum ⁽⁴⁾ (East of Rocky Mountains)	two-spotted spider Banks grass	X	1½ to 2 pints	minimum of 20 gallons	minimum of 5 gallons	1	30 ⁽⁵⁾
Potatoes (Pacific Northwest Only)	two-spotted spider	X	1½ to 2½ pints ⁽⁶⁾	20 to 50 gallons	minimum of 10 gallons	2	14
Beans (green lima and dry) (West of Rocky Mountains)	two-spotted spider	X	2-3 pints	minimum of 20 gallons	minimum of 5 gallons	2	7 ^(7,8)
Oranges and Grapefruit (Florida and Texas only)	citrus red Texas citrus six-spotted citrus rust false spider	X	30-120 Fluid Ounces ⁽⁹⁾	600-1500 gallons	X	2	7
			2-3 pints	minimum of 25 gallons	minimum of 10 gallons		

APPLICATION NOTES

COTTON

1. COMITE may be phytotoxic on young cotton under certain conditions. Consult your Uniroyal representative before spraying cotton less than 10 inches in height.
2. COMITE may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

CORN

3. Includes grain and silage.

GRAIN SORGHUM

4. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check for phytotoxicity before using COMITE on a large area.
5. SILAGE STAGE. Grain should not be harvested within 60 days of spraying COMITE. Do not plant rotational crops within 6 months of the last treatment on grain sorghum.

POTATOES

6. The 2½ pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

7. Succulent beans. Days before harvest (last spray) for dry beans is 28 days. Do not plant rotational crops within 6 months of the last treatment on beans.
8. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

ORANGES AND GRAPEFRUIT

9. Amount of COMITE per 500 Gallon Tank — 25-40 Fluid Ounces.

CAUTION!

On citrus, leaf injury may occur if COMITE is tank mixed with oil or applied within two weeks prior to or following an oil application.

2/2