

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
DANGER**

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

3. CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

4. GENERAL

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

COMITE®

AGRICULTURAL MITICIDE

UNIROYAL

COMPOSITION

Active Ingredient: (% by weight)	
Propargite 2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite*	73.0%
Inert Ingredients:	27.0%
Total:	100.0%

*U.S. Patents 3,272,854 - 3,463,859

**Contains 6.55 pounds OMITE per gallon.

**KEEP OUT OF REACH OF CHILDREN.
DANGER**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-104-ZA

NET WEIGHT

11-16-81

A liquid emulsifiable concentrate for preparing sprays to control the Banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus, and two-spotted spider mites.

COMITE is not systemic in action, therefore, complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of COMITE to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

See back side for dosage instructions chart and application notes.

AL USUARIO:

Si usted no puede leer o entender Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TO THE USER:

If you cannot read or understand English, do not use this product until the label has been fully explained to you.

USE RESTRICTIONS —

Cotton — Apply only before bolls open. Do not feed treated foliage or cotton trash to live-stock.

Beans — Do not feed or forage treated bean vines or bean trash after harvesting.

Peanuts — Do not graze or feed livestock on treated areas or cut treated forage or hay.

Do not graze or feed livestock on cover crops grown among the trees listed on this label.

General — Do not mix COMITE with alkaline materials (such as lime, Bordeaux mixture and lime sulfur) or with materials containing a large amount of petroleum type solvents

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

1974 UNIROYAL Inc All Rights Reserved

DOSAGE INSTRUCTIONS

Crop	Mites Controlled	Timing of Application	Amount of COMITE Per Acre	Gallons of Spray Solution Per Acre		Total Number of Sprays Per Year	Days Before Harvest (Last Spray)
				Ground Application	Aerial Application		
Cotton	two-spotted, Pacific and strawberry spider	Early (1)	1 pint	15 (2) to 30 gallons	Not recommended	3	Before bolls open
		Mid Season to Layby	1 1/2 to 2 pints	25 (2) to 40 gallons	—		
			2 pints	—	5 to 15 gallons		
		Layby to Boll Opening	2 pints	25 (2) to 50 gallons	5 to 15 gallons		
Field corn (3-4) (grain & silage)	two-spotted spider, Banks grass	X	2-3 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
Grain Sorghum (5) (East of Rocky Mountains)	two-spotted spider, Banks grass	X	1 1/2 to 2 pints	minimum of 20 gallons	minimum of 5 gallons	1	30
Peanuts	two spotted spider	X	2 pints	minimum of 20 gallons	minimum of 5 gallons	1	14
Potatoes (Pacific Northwest Only)	two spotted spider	X	1 1/2 to 2 1/2 pints (7)	20 to 50 gallons	minimum of 10 gallons	2	14
Beans (6) (green lima and dry) (West of Rocky Mountains)	two spotted spider	X	2-3 pints	minimum of 20 gallons	minimum of 5 gallons	2	succulent: 7 dry: 28
Oranges and Grapefruit (Florida and Texas only)	citrus red, Texas citrus six spotted, citrus rust, false spider	X	30-120 Fluid Ounces (5)	600-1500 gallons	X	2	7
			2-3 pints	minimum of 25 gallons	minimum of 10 gallons		

← INSERT A

APPLICATION NOTES

COTTON

1. COMITE may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
2. COMITE may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE dosage rate as listed.

CORN

3. Treat early, best results are achieved when mite populations are low, before the mites have damaged the plants. We recommend treatment when mite populations are just beginning to build and the corn is still small enough (usually 3 to 4 feet) to allow good coverage.
4. In Kansas, Colorado and New Mexico use a minimum of 2 gallons spray solution per acre. For all other states use a minimum of 5 gallons spray solution per acre.

GRAIN SORGHUM

5. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE on a large area.
6. SILAGE STAGE Grain should not be harvested within 60 days of spraying COMITE.

POTATOES

7. The 2 1/2 pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

8. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

ORANGES AND GRAPEFRUIT

9. Amount of COMITE per 500 Gallon Tank — 25-40 Fluid Ounces

CAUTION

On citrus, leaf injury may occur if COMITE is tank mixed with oil or applied within two weeks prior to or following an oil application.

INSERT A

Sweet Corn ⁽³⁾ (grain & silage) (California only)	Two-spotted spider Banks grass	X	2-3 pints	Minimum of 10 gallons	Minimum of 1 gallons	1	30
--	--------------------------------------	---	-----------	-----------------------------	----------------------------	---	----

355/17382
18/16

TRP
1/12/83

Mr. Bogislav von Schmeling
Uniroyal Chemical
Division of Uniroyal Inc.
Crop Protection, Research and Development
74 Aaltje Road
Bethany, CT 06525

Dear Mr. Von Schmeling:

Subject: Amendment-Add Sweet Corn - California Only
COMITE
EPA Registration No. 400-104
Your Application Dated June 14, 1982

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The basis for the acceptance of this added use on sweet corn in California at the rate of 2-3 pints with a 30-day Pre-Harvest Interval was examination of sweet corn residue data with PP #042424 and the assumption that residues of propargite will not be different than those in field corn forage and fodder from the same use.