

COMITE

COMITE®

AGRICULTURAL MITICIDE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

DANGER

Corrosive, causes eye damage. May be fatal if inhaled. Harmful if swallowed or absorbed through skin. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wear goggles or face shield when handling. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and shrimp. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

UNIROYAL

ACCEPTED
JAN 21 1986
EPA Reg. No. 400-104

COMPOSITION

Active Ingredients: (% by weight)	
Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]*	73.0%
Inert Ingredients:	27.0%
Total:	100.0%

*U.S. Patent No. 3,463,859

**Contains 6.55 pounds OMITE per gallon

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

See left side panel for additional precautionary statement.

UNIROYAL CHEMICAL COMPANY, INC.

Middlebury, CT 06749

EPA REG. NO. 400-104

EPA EST. NO.

Lot. No.

11-5-85

COMITE is a Registered Trademark of UNIROYAL CHEMICAL COMPANY, INC.

©Copyright 1984 Uniroyal Chemical Company, Inc.

NET WEIGHT:

REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to the workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER, Area to be treated with Propargite on [date of application]. Do not enter without appropriate protective clothing until sprays have dried. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid recommendations."

GENERAL INSTRUCTIONS

COMITE is a liquid emulsifiable concentrate for preparing sprays to control the banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus, and two-spotted spider mites.

COMITE is not systemic in action; therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of COMITE to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	TIMING OF APPLICATION	AMOUNT OF COMITE PER ACRE	GALLONS OF SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
				Ground Application	Aerial Application		
COTTON (California & Arizona only)	two-spotted spider Pacific spider strawberry spider	Early ⁽¹⁾	1 pint	15 ⁽²⁾ to 30 gallons	Not recommended	3	Before bolls open
		Mid Season to Layby	1½ to 2 pints	25 ⁽²⁾ to 40 gallons	—		
			2 pints	—	5 to 15 gallons		
		Layby	2 pints	25 ⁽²⁾ to 50 gallons	5 to 15 gallons		
		Layby Boll Opening	2 pints	—	5 to 15 gallons		
COTTON (East of the Rocky Mts.)	two-spotted spider Pacific spider Strawberry spider	Early ⁽¹⁾	1 pint	15 ⁽²⁾ to 30 gallons	Not recommended	3	Before bolls open
		Mid Season to Layby	1 to 2 pints	25 ⁽²⁾ to 40 gallons	5 to 15 gallons		
			Layby to Boll Opening	2 pints	25 ⁽²⁾ to 50 gallons		
FIELD CORN ⁽³⁾	two-spotted spider Banks grass	 	2 to 3 pints	minimum of 20 gallons	minimum of 2 gallons ⁽⁴⁾	1	30
SWEET CORN ⁽³⁾ (California only)	two-spotted spider Banks grass	 	2 to 3 pints	minimum of 20 gallons	minimum of 2 gallons	1	30
GRAIN SORGHUM ⁽⁵⁾ (East of Rocky Mountains)	two-spotted spider Bank grass	 	1½ to 2 pints	minimum of 20 gallons	minimum of 5 gallons	1	30 ⁽⁶⁾
PEANUTS	two-spotted spider	 	2 pints	minimum of 20 gallons	minimum of 5 gallons	1	14
POTATOES (Pacific North-west Only)	two-spotted spider	 	1½ to 2½ pints ⁽⁷⁾	20 to 50 gallons	minimum of 10 gallons	2	14
BEANS ⁽⁸⁾ (green, lima & dry) (West of Rocky Mts.)	two-spotted spider	 	2 to 3 pints	minimum of 20 gallons	minimum of 5 gallons	2	succulent - 7 dry - 28
ORANGES & GRAPEFRUIT ⁽⁹⁾ (Florida and Texas only)	citrus red Texas citrus six-spotted citrus rust false spider broad	 	2 to 3 pints	minimum of 25 gallons	minimum of 10 gallons	2	7

() See respective application note on side panel.

14

APPLICATION NOTES

COTTON

1. COMITE may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
2. COMITE may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE dosage-rate per acre.

CORN

3. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. In California, ensure applications are made only when corn leaves are dry. This is especially true for white corn varieties which are the most susceptible to plant injury.
4. In Kansas and Colorado, use a minimum of 2 gallons spray solution per acre. In California, best results are obtained from applying 10-20 gallons of spray solution per acre. In Texas and New Mexico and other states, use a minimum of 5 gallons spray solution per acre.

GRAIN SORGHUM

5. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE on a large area.
6. SILAGE STAGE: Grain should not be harvested within 60 days of spraying COMITE.

POTATOES

7. The $\frac{1}{2}$ pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

8. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

ORANGES AND GRAPEFRUIT

9. Apply in sufficient water for thorough coverage.

On citrus, leaf distortion and/or fruit spotting may occur when used in the spring or it may occur if COMITE is tank mixed with oil or applied within two weeks prior to or following an oil application.

Do not use in spray solution above pH 10.

USE RESTRICTIONS

Cotton - Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.

Beans - Do not feed or forage treated bean vines or bean trash after harvesting.

Peanuts - Do not graze or feed livestock on treated areas or cut treated forage for hay.

Do not graze or feed livestock on cover crops grown among the trees listed on this label.

General - Do not mix COMITE with alkaline materials (such as lime, Bordeaux mixture or lime sulfur) or with materials containing a large amount of petroleum type solvents.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.