

DANGER! KEEP OUT OF REACH OF CHILDREN.

Do not get in eyes.
 Causes severe eye damage.
 Wear goggles when handling.
 In case of contact with eyes, flush immediately with running water and get medical attention.
 For other statements see right panel.

VITAVAX®-3F FLOWABLE FUNGICIDE

COLLATERAL

NET CONTENTS


COMPOSITION

ACTIVE INGREDIENT (% by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathiazin-3-carboxamide)**	34.0%
INERT INGREDIENTS	66.0%
TOTAL	100.0%

*U.S. Patents 3,249,455 - 3,593,202 - 3,454,391
 **Contains 3.34 pounds of active Carboxin per gallon.

EPA Reg. No. 400-107-AA

EPA Est. No.



DIRECTIONS FOR USE

VITAVAX-3F is a ready to use seed treatment which should be applied as a slurry treatment by machine.

BARLEY AND WHEAT

For the control of loose smut (*Ustilago nuda*), covered smut (*Ustilago hordei*) of barley, loose smut (*Ustilago tritici*) and common bunt (*Tilletia caries*) of wheat. Use at the rate of 2-3 fluid ounces per 100 pounds of seed.

USE RESTRICTIONS

Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

COMPATIBILITY

VITAVAX-3F can be used on seed previously treated or to be treated at the recommended rate of other seed treatment fungicides such as thiram.

HOWEVER, VITAVAX-3F SHOULD NOT BE TANK MIXED WITH ANY PESTICIDE BECAUSE OF POSSIBLE PHYSICAL INCOMPATIBILITIES BETWEEN FORMULATIONS.

CORN

VITAVAX-3F controls seed rots and provides systemic control of seedling diseases of corn. VITAVAX-3F is particularly effective against *Rhizoctonia solani*. VITAVAX-3F should be used at a rate of 2-4 fluid ounces per 100 pounds of seed when applied to seed that has previously been treated or will be treated with Protectant Fungicides such as captan or thiram.

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

COMPATIBILITY

VITAVAX-3F can be used on seed previously treated or to be treated with the recommended rate of other seed treatment fungicides such as captan or thiram. HOWEVER, VITAVAX-3F SHOULD NOT BE TANK MIXED WITH ANY PESTICIDE BECAUSE OF POSSIBLE PHYSICAL INCOMPATIBILITY BETWEEN FORMULATIONS.

DANGER

HANDLING — Harmful if swallowed. Avoid inhaling spray. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Keep away from food or food products. Care must be exercised in the handling of treated seed.

SEED LABELING — "This seed treated with VITAVAX-3F. Do not use for food, feed or oil purposes. On corn, barley, and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments.

COMPATIBILITIES — Observe all cautions and limitations on labeling of all products used in mixture with this product. Apply this product only as specified on this label.

FISH AND WILDLIFE — Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL — Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food or water. If disposal is by burning, keep out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

11-23-76-5FF

Lot No.
 © 1973 Uniroyal, Inc. All Rights Reserved.

DANGER! KEEP OUT OF REACH OF CHILDREN.

Do not get in eyes.

Causes severe eye damage.

Wear goggles when handling.

In case of contact with eyes, flush immediately with running water and get medical attention.

For other statements see right panel

VITAVAX[®]-3F FLOWABLE FUNGICIDE

BASIC

NET CONTENTS

COMPOSITION

ACTIVE INGREDIENT (% by weight)	
Carboxin (5,6-d.h., dro-2-methyl-1,4-oxathio-2-carboxanilide)*	34.0%**
INERT INGREDIENTS	66.0%
TOTAL	100.0%

*U.S. Patents 3,249,499 - 3,393,222 - 3,454,351

**Contains 3.34 pounds of active Carboxin per gallon.

EPA Reg. No. 400-107-AA

EPA Est. No.

ACCEPTED

DEC 7 1976

U.S. Department of Agriculture, Act, registered under EPA Reg. No. 400-107

UNIROYAL

DIRECTIONS FOR USE

VITAVAX-3F is a ready to use seed treatment which should be applied as a slurry treatment by machine.

BARLEY AND WHEAT

For the control of loose smut (*Ustilago nuda*), covered smut (*Ustilago hordei*) of barley, loose smut (*Ustilago tritici*) and common bunt (*Tilletia caries*) of wheat. Use at the rate of 2-3 fluid ounces per 100 pounds of seed.

USE RESTRICTIONS

Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

COMPATIBILITY

VITAVAX-3F can be used on seed previously treated or to be treated at the recommended rate of other seed treatment fungicides such as thiram.

HOWEVER, VITAVAX-3F SHOULD NOT BE TANK MIXED WITH ANY PESTICIDE BECAUSE OF POSSIBLE PHYSICAL INCOMPATIBILITIES BETWEEN FORMULATIONS.

COTTON

VITAVAX-3F controls seed rots and provides systemic control of seedling diseases of cotton. Seed rot, damping-off and sore-shin due to *Rhizoctonia solani* are prevented. VITAVAX-3F may be used on both machine and acid delinted cottonseed. VITAVAX-3F should be used at the rate of 8 fluid ounces per 100 pounds of seed.

USE RESTRICTIONS

Do not use treated cottonseed for food, feed or oil purposes. Do not graze or feed livestock on hay grown from treated seed.

COMPATIBILITY

VITAVAX-3F can be used on seed previously treated or to be treated at the recommended rate of other seed treatment fungicides, such as captan or thiram, or seed treatment insecticides, such as Disyston or Thimet.

HOWEVER, VITAVAX-3F SHOULD NOT BE TANK MIXED WITH ANY PESTICIDE BECAUSE OF POSSIBLE PHYSICAL INCOMPATIBILITY BETWEEN FORMULATIONS.

CORN

VITAVAX-3F controls seed rots and provides systemic control of seedling diseases of corn. VITAVAX-3F is particularly effective against *Rhizoctonia solani*. VITAVAX-3F should be used at a rate of 2-4 fluid ounces per 100 pounds of seed when applied to seed that has previously been treated or will be treated with Protectant Fungicides such as captan or

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

COMPATIBILITY

VITAVAX-3F can be used on seed previously treated or to be treated with the recommended rate of other seed treatment fungicides such as captan or thiram. HOWEVER, VITAVAX-3F SHOULD NOT BE TANK MIXED WITH ANY PESTICIDE BECAUSE OF POSSIBLE PHYSICAL INCOMPATIBILITY BETWEEN FORMULATIONS.

DANGER

HANDLING — Harmful if swallowed. Avoid inhaling spray. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Keep away from food or food products. Care must be exercised in the handling of treated seed.

SEED LABELING — "This seed treated with VITAVAX-3F. Do not use for food, feed or oil purposes. On corn, barley, and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments.

COMPATIBILITIES — Observe all cautions and limitations on labeling of all products used in mixture with this product. Apply this product only as specified on this label.

FISH AND WILDLIFE — Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL — Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food or water. If disposal is by burning, keep out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

11-22-76-5FF

Lot No.