

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS  
CAUTION**

AVOID CONTACT WITH SKIN, EYES OR CLOTHING. WASH HANDS AND FACE THOROUGHLY WITH SOAP AND WATER AFTER USE AND BEFORE EATING OR SMOKING.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all treated seeds by burial away from bodies of water.

**PHYSICAL or CHEMICAL HAZARDS**  
Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

For use directions, see label on container. Do not use this product in a manner inconsistent with the label.

**STORAGE AND DISPOSAL**

1. PROHIBITIONS
2. PESTICIDE DISPOSAL
3. CONTAINER DISPOSAL
4. GENERAL

ACCORD  
 OCT 12 1978  
 U.S.P.  
 22 Oct  
 112-167

**VITAVAX<sup>®</sup>**  
FLOWABLE FUNGICIDE



**COMPOSITION**

Active ingredient (by weight):  
 Carboxin (5,6-dihydro-2-methyl-4-oxathin-3-carboxamide)  
 Inert ingredient  
 Total  
 U.S. Patents 3,249,499; 3,393,791; 3,414,477  
 \*\*Contains 3.34 pounds of active Carboxin per gallon.

**KEEP OUT OF REACH OF CHILDREN.  
CAUTION!**

**STATEMENTS OF PRACTICAL TREATMENT**

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled: See a physician if an abnormal reaction occurs.

If on skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water and get medical treatment.

See left side panel for additional precautionary statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, INC.  
 Naugatuck, Connecticut 06770  
 EPA Establishment No.  
 EPA Registration No. 400-107-AA

NET WEIGHT

9 20 78

**DIRECTIONS FOR USE**

VITAVAX Flowable Fungicide is a ready to use seed treatment which should be applied as a slurry treatment by machine.

**BARLEY AND WHEAT**

For the control of loose smut (Ustilago nuda), covered smut (Ustilago hordei) of barley, loose smut (Ustilago tritici) and common bunt (Tilletia caries) of wheat. Use at the rate of 2-4 fluid ounces per 100 pounds of seed.

**CORN**

VITAVAX Flowable Fungicide controls seed-bots and provides systemic control of seedling diseases of corn. VITAVAX Flowable Fungicide is particularly effective against *Peridermium solani*. VITAVAX Flowable Fungicide should be used at a rate of 2-4 fluid ounces per 100 pounds of seed when applied to seed that has previously been treated or will be treated with protectant fungicides such as captan or thiram.

**COMPATIBILITY**

VITAVAX Flowable Fungicide can be used on seed previously treated or to be treated with the recommended rate of other seed treatment fungicides such as captan or thiram.

**FOR CORN SEED ONLY**

VITAVAX Flowable Fungicide may be tank mixed with the following registered pesticides: Orthocide<sup>®</sup> 75 Seed Protectant, Orthocide<sup>®</sup> Methoxychlor 75-3 Seed Protectant, Captan<sup>®</sup> Seed Protectant, Captan Methoxychlor 75-2 Seed Protectant, and Cythion<sup>®</sup> 8-F. Do not allow mixtures to stand overnight. Thoroughly clean equipment between treating operations.

Orthocide<sup>®</sup> trademark of Chevron Chemical Company.

Cythion<sup>®</sup> trademark of American Cyanamid.

Observe all cautions and limitations on labeling of all products used in mixture with this product only as specified on this label.

#### USE RESTRICTIONS

When using formulations that do not contain dye, to comply with 21CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180.1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals. Treated seed must not be used for or mixed with food of animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Do not graze or feed livestock on treated areas for six weeks after planting.

#### SEED LABELING

"Treated seed must not be used for or mixed with food of animal feed, or processed for oil. Seed commercially treated with VITAVAX Flowable Fungicide must be labeled in accord with all applicable requirements of the Federal Seed Act. On corn, barley and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments.

**IMPORTANT NOTICE** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the direction, and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose express or implied extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.