

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
CAUTION**

AVOID CONTACT WITH SKIN, EYES
OR CLOTHING. WASH HANDS AND
FACE THOROUGHLY WITH SOAP AND
WATER AFTER USE AND BEFORE
EATING OR SMOKING.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply
directly to lakes, ponds or streams. Do not
contaminate water by cleaning of equip-
ment or disposal of wastes.

Exposed treated seeds may be hazardous
to birds and other wildlife. Dispose of all
treated seeds by burial away from bodies of
water.

PHYSICAL or CHEMICAL HAZARDS
Do not use or store near heat or open flame.

DIRECTIONS FOR USE

For use in accordance with Federal Law. Use this
product only on the crops and weeds listed on the label.
Do not use on other crops.

STORAGE AND DISPOSAL

1. PROHIBITIONS

Do not contaminate water, food, or
feed by storage and disposal. Open
dumping is prohibited.

2. PESTICIDE DISPOSAL

Pesticide spray, wash, or residue
that cannot be used on chemicals, re-
siduals should be disposed of in a
safe, approved manner. Do not
dump in a safe place away from water
supply.

3. CONTAINER DISPOSAL

Triple rinse or equivalent and offer to
recycling or disposal, approved and
labeled for use in a safe place.

4. GENERAL

Consult Federal, State and Local
Health Authorities for approved dis-
posal methods. Do not use in a
manner inconsistent with the label.



VITAVAX®

FLOWABLE FUNGICIDE

UNIROYAL

COMPOSITION

Active ingredient: Captan 80 34.0%
Inert ingredient: 66.0%
Total: 100.0%
Net weight: 100.0%
EPA Reg. No. 400-107 AA

KEEP OUT OF REACH OF CHILDREN.

CAUTION!

STATEMENTS OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of
throat with finger or blunt object. Do not induce vomiting or give anything by
mouth to an unconscious person. Get medical attention.

If inhaled: See a physician if an abnormal reaction occurs.

If on skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water and get medical treatment.

See left side panel for additional precautionary statement.

Mfg. by UNIROYAL CHEMICAL, Division of UNIROYAL, INC.
Naugatuck, Connecticut 06770
EPA Establishment No.
EPA Registrat. No. 400-107 AA

NET WEIGHT

1-28-81

DIRECTIONS FOR USE

VITAVAX Flowable Fungicide is a ready to
use seed treatment which should be ap-
plied as a dry treatment by machine.

BARLEY, WHEAT AND OATS

For the control of loose smut (*Ustilago
nuda*), covered smut (*Ustilago hordei*) of
barley; loose smut (*Ustilago avenae*),
covered smut (*Ustilago kolleri*) of oats;
loose smut (*Ustilago tritici*) and common
bunt (*Tilletia caries*) of wheat. Use at the
rate of 2-3 ounces per 100 pounds of seed.

CORN

VITAVAX FLOWABLE FUNGICIDE con-
trols seed and seedling diseases and
provides systemic control of seedborne
head smut (*Sphacelotheca reiliana*).
VITAVAX FLOWABLE FUNGICIDE is
also particularly effective against *Rhizoc-
tonia solani*. VITAVAX FLOWABLE
FUNGICIDE should be used at a rate of 2
to 4 fluid ounces per 100 pounds of seed
when applied to seed that has been treated
or will be treated with protectant
fungicides such as captan or thiram.

COMPATIBILITY

VITAVAX Flowable Fungicide can be used
in combination with Gustafson's CAPTAN,
EPA Reg. No. 400-107 AA, or Gustafson's
4-5 Flowable seed protectants. All other
tank mixes should be pretested to
determine physical compatibility between
formulations, and to assure there will be no
adverse phytotoxic reaction.

FOR CORN SEED ONLY

VITAVAX Flowable Fungicide may be tank
mixed with the following registered prod-
ucts: Orthocide® 75-3 Seed Protectant,
Orthocide® Methoxychlor 75-3 Seed Pro-
tectant, Captan 80 Seed Protectant, Captan
Methoxychlor 75-2 Seed Protectant, and
Cythion® 4-F. Do not allow mixtures to
stand overnight. Thoroughly clean equip-
ment between treating operations.

Orthocide® is a trademark of Chevron
Chemical Company.

Cythion® is a trademark of American Cyan-
amid.

See label, caution and limitations on
labeling of all product used in mixture
with this product only as specified on the
label.

When using a granulator, granulate the seed with dye to comply with the Federal Seed Act, Section 125. Do not granulate seed with dye if the granulator has not been tested and proved dye 4000 R for 1000000 seeds and prevent subsequent and subsequent use as a food for man or feed for animal. Treated seed must not be used to be mixed with food of animal feed, or processed for oil. Treat only those seed needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Do not graze or feed livestock on treated areas for six weeks after planting.

SEED LABELING

"Treated seed must not be used for or mixed with food of animal feed, or processed for oil. Seed commercially treated with VITAVAX Flowable Fungicide must be labeled in accord with all applicable requirements of the Federal Seed Act. On corn, barley, and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.