

DIRECTIONS FOR USE

VITAVAX Flowable Fungicide is a ready to-use seed treatment which should be applied as a slurry treatment by machine.

BARLEY, WHEAT AND OATS

For the control of loose smut (Ustilago nuda), covered smut (Ustilago hordel), of barley, Hoose smut (Ustilago avenae), covered smut (Ustilago kolleri) of oats, loose smut (Ustilago tritici) and common bunt (Tilletia caries) of wheat. Use at the rate of 2-3 ounces per 100 pounds of seed.

CORN

VITAVAX Flowable Fungicide controls seed and seedling diseases and provides systemic control of seedborne head smut (**Sphacelotheca reiliana**). VITA-VAX Flowable Fungicide is also particularly effective against **Rhizoctonia solani**. VITAVAX Flowable Fungicide should be used at a rate of 2 to 4 ounces per 100 pounds of seed when applied to seed that has been treated or will be treated with protectant fungicides such as captan or thiram.

COTTON

VITAVAX Fungicide controls seed rots and provides systemic control of seedling diseases of cotton. Seed rot, damping off and sore-shin due to **Rhizoctonia** solani are prevented. VITAVAX may be used on both machine and acid delinted cottonseed. Use 3-8 ounces of VITAVAX per 100 pounds of seed.

VITAVAX Flowable Fungicide can be used on seed previously treated or to be treated with other seed treatment fungicides, such as captan or thiram, or seed treatment insecticides, such as Disyston or Thimet.

COMPATIBILITY

VITAVAX Flowable Fungicide can be used in combination with Gustafson,s CAPTAN 30-DD and CAPTAN-300 or Gustafson's 42-S flowable seed protectants. All other tank mixes should be pre-tested to determine physical compatibility between formulations, and to assure there will be no adverse phytotoxic reactions.

FOR CORN SEED ONLY

VITAVAX Flowable Fungicitie may be tank mixed with the following tegis tered pesticides: Orthocide: 75 Seed Protectant, Orthocide: Methoxychlor 75-3 Seed Protectant, Captan 80 Seed Protectant, Captan Methoxychlor 75-2 Seed Protectant, and Cythion** 8-F Do not allow mixtures to stand overnight. Thoroughly clean equipment between treating operations.

*Orthocide® trademark of Chevron Chemical Company.

**Cythion® trademark of American Cyanamid.

Observe all cautions and limitations on labeling of all products used in mixture with this product only as specified on this label.

USE RESTRICTIONS

When using formulations that do not contain dye, to comply with 21CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 108.1001) to distinguish and prevent subsequent and inadvertent use as food for man or feed for animals. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Do not graze or feed livestock on treated areas for six weeks after planting.

SEED LABELING

"Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Seed commercially treated with VITAVAX Flowable Fungicide must be labeled in accord with all applicable requirements of the Federal Seed Act. On corn, barley, and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments. IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warfanty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

 $^{\odot}$ 1973 Uniroyal, Inc. All Rights Reserved.