

DANGER! KEEP OUT OF REACH OF CHILDREN.

Do not get in eyes.
Causes severe eye damage.
Wear goggles when handling.
In case of contact with eyes, flush immediately with running water and get medical attention.
For other danger statements see right panel.

VITAVAX-200

FLOWABLE FUNGICIDE

(VITAVAX WITH THIRAM)

Net Contents 55 LIQUID GALLONS

May 11, 1974

400-112

COMPOSITION	
Active Ingredients: (by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathin-3-carboxanilide) *	17.0
Thiram (tetramethylthiuram disulfide)	17.0
Inert Ingredients	66.0
Total	100.0%

* U.S. Patents 3,249,499; 3,393,202; 3,454,391
EPA Reg. No. 400-112-AA



DIRECTIONS FOR USE

VITAVAX-200 Flowable Fungicide is a ready-to-use seed treatment.

VITAVAX-200 Flowable Fungicide combines the systemic action of carboxin with the surface action of thiram to control various seed and seedling diseases of barley and wheat. It is particularly effective against Rhizoctonia solani.

In addition, loose smut (Ustilago nuda) and covered smut (Ustilago hordei) are controlled on barley, and loose smut (Ustilago tritici) and common bunt (Tilletia caries) are controlled on wheat by VITAVAX-200 Flowable Fungicide.

Barley and Wheat

Use 3-4 fluid ounces of VITAVAX-200 Flowable Fungicide per 100 pounds of seed. VITAVAX-200 Flowable Fungicide should be applied as a slurry treatment by machine.

Use Restrictions

Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting.

Compatibility

VITAVAX-200 Flowable Fungicide can be used in combination with the commonly used barley and wheat seed protectants. However, VITAVAX-200 Flowable Fungicide cannot be directly mixed with fungicides or insecticides prior to application.

Lot No. _____

© 1972 UNIROYAL, Inc. All Rights Reserved

UNIROYAL Chemical - Division of UNIROYAL, Inc. - Naugatuck, Connecticut 06770

DANGER

HANDLING — Harmful if swallowed. Avoid inhaling spray. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Keep away from food or food products. Care must be exercised in the handling of treated seed. Consumption of alcoholic beverages increases the possibility of harm.

SEED LABELING — "This seed treated with VITAVAX-200 Flowable Fungicide. Do not use treated seed for food, feed or oil purposes." Add appropriate information for any other chemical treatments.

FISH and WILDLIFE — This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONTAINER DISPOSAL — Bury container 18 inches below soil surface at safe disposal site. Never reuse for food or water.

IMPORTANT NOTICE Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

5-74-55G