

ACCEPTED

SEP 22 1983

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of registration of this pesticide.

EPA Reg. No. 400-112

VITAVAX-200

FLOWABLE FUNGICIDE
(VITAVAX WITH THIRAM)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Consumption of alcoholic beverages increases the possibility of harm.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

UNIROYAL

COMPOSITION

Active Ingredients (by weight):	
Carboxin (5,6-dihydro-2-methyl-1,4-dioxathin-3-carboxanilide)*	17.0
Thiram (tetramethylthiuram disulfide)	17.0
Inert Ingredients	66.0
Total	100.0

*U.S. Patents 3,249,499 - 3,393,202 - 3,454,391

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and see a physician.
IF INHALED: See a physician if an abnormal reaction occurs.
IF ON SKIN: Wash thoroughly with soap and water.
IF IN EYES: Immediately flush eyes with plenty of water. See a physician.
SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL Inc.
Naugatuck, Connecticut 06770
EPA REG NO 400 112
EPA EST NO
VITAVAX is a Registered Trademark of UNIROYAL Inc.
Copyright 1983, Uniroyal Inc.

8 19 83

L I No

NET WEIGHT

DIRECTIONS FOR USE

VITAVAX-200 Flowable Fungicide is a ready-to-use seed treatment

RICE

VITAVAX-200 Flowable Fungicide combines the systemic action of carboxin with the surface action of thiram to control various seed and seedling diseases of rice. It is particularly effective against *Rhizoctonia solani* and *helminthosporium oryzae*.

Use 3-4 fluid ounces of VITAVAX-200 Flowable Fungicide per 100 pounds of seed applied as a slurry treatment by machine. The higher the rate is recommended for control of *Helminthosporium oryzae*.

BARLEY, WHEAT AND OATS

In addition, loose smut (*Ustilago nuda*), and covered smut (*Ustilago hordei*), are controlled on barley and loose smut (*Ustilago tritici*), and common bunt (*Tilletia caries*) are controlled on wheat by VITAVAX-200 Flowable Fungicide.

Use 3-4 fluid ounces of VITAVAX-200 Flowable Fungicide per 100 pounds of seed. VITAVAX-200 Flowable Fungicide should be applied as a slurry treatment by machine.

SOYBEANS

Use 2 fluid ounces of VITAVAX-200 Flowable Fungicide per bushel of seed or 4 fluid ounces per 100 pounds of seed. VITAVAX-200 Flowable Fungicide should be applied as a slurry treatment by machine or to the hopper box as a pour on treatment.

USE RESTRICTIONS

Do not use treated seed for food or feed purposes. Do not graze or feed livestock on treated areas for six weeks after planting. Care must be exercised in the handling of treated seed. Do not graze or feed livestock on vines grown from treated seed.

USE PRECAUTION

When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180.1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals.

COMPATIBILITY

VITAVAX-200 Flowable Fungicide should not be tank mixed with any pesticide unless adequate physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.

SEED LABELING

"This seed treated with VITAVAX-200 Flowable Fungicide. Do not use treated seed for food, feed, or oil purposes. Add appropriate information for any other chemical treatments."

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.