400-113 PM21 1022

VITAVAX®-17



as amenawa tegistered under EPA Reg. No.

ACCEPTED

JUN 2 1986 Under the Federal Insecticide.

Fungleide, and Rodenticide Act. as amended, for the pesticide

Causes eye irritation. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothiny. Wash hands and face thoroughly with soap and water after use and before eating or smoking.

ENVIRONMENTAL HAZARDS Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

ſ

PHYSICAL or CHEMICAL HAZARDS Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

UNIROYAL

**Contains 1.50 pounds of active Carboxin per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching basis of throat with linger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: See a physician II an abnormal reaction occurs .

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately Ilush eyes with plenty of water and get medical treatment. SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT...

Uniroyal Chemical Company, Inc. Middlebury, CT 06749

EPA REG. NO. 400-113-AA EPA EST. NO. VITAVAX is a Registered Trademark of UNIROYAL, Inc. Copyright 1983. Uniroyal, Inc.

Loi. No.

• 4-14-86

. .

NET WEIGHT:

2062

DIRECTIONS FOR USE

is a ready-to-use-seed treatment which should be applied as a slurry treatment VITAVAX - 17 by machine.

BARLEY, DATS, WHEAT AND TRITICALE

For the control of loose smut (Usiliago nuda), covered smut (Usiliago hordei), of barley, loose smut (Usiliago avenael, covered smut (Ustilago kolleri), of oats, loose smut (Ustilago tritici) and common bunt (Tilletla caries) of wheat. Use at the rate of <u>ounces</u> per 100 pounds of seed.

(4-6 f]CORN controls seed and seedling diseases and provides systemic control of seedborne VITAVAX -17 head smut (Sphacelotheca reiliana). VITAVAX - 17 is also particularly effective against Rhizoctonia solani. VITAVAX Flowable Fungicide should be used at a rate of _____ ounces per 100 pounds of seed when applied to seed that has been treated or will be treated with protectant/lungicides such as captan or thiram.

(4-8 f1)COTTON

VITAVAX-17 Flowable Fungicide provides systemic control of seed and seedling diseases of cotton. Seed rot, damping off and sore-shin due to Rhizoctonia solani are prevented. VITAVAX-17 Flowable Fungicide ct 5e used on both machine and acid delinted cottonseed at the rate of 16 fluid ounces per 100 pounds of seed.

BEANS

should be used at the rate of 6-8 fluid ounces per 100 pounds of seed for the or beans. VITAVAX -17 control of seedling diseases for beans. Do not allow livestock to graze or feed on bean forage until 60 days after treated seed is planted.

- SOYBEANS

VITAVAX -17 \cdot should be used at the rate of \mathcal{N} fluid ounces per 100 pounds of seed for the control of seedling diseases in soybeans. 6-8

Do not graze or feed livestock on forage and hay on treated areas.

COMPATIBILITY

VITAVAX - 17 can be used in combination with Gustafson's Captan 30-DD and CAPTAN-300 or Gustafson's 42-S flowable seed protectants. All other tank mixes should be pre-tested to determine physical compatibility between formulations, and to assure there will be no adverse phytotoxic reactions.

FOR CORN SEED ONLY

VITAVAX -17 . may be tank mixed with the following registered pesticides: Orthocide* 75 Seed Protectant, Orthocide* Methoxychlor 75-3 Seed Protectant, Captan 80 Seed Protectant, Captan Methoxychlor 75-2 Seed Protectant, and Cythion** 8-F. Do not allc mixtures to stand overnight. Thoroughly clean equipment between treating operations.

*Orthocide® trademark of Chevron Chemical Company.

**Cythion® trademark of American Cyanamid.

Observe all cautions and limitations on labeling of all products used in mixture with this product only as (specified on this label.

USE RESTRICTIONS

This product contains a dye which imparts an unnatural color to the seed to help prevent the inadvertant use of treated seed for oil purposes or, as foud for man or feed for animals. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, ٠. e t minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting ****

time. Do not graze or feed livestock on treated areas for six weeks after planting.

SEED LABELING

"Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Seed commercially treated with WJAVAX Flowable Fungicide must be labeled in accord with all applicable requirements of the Federal Seed Act. On corn. barley. and wheat, do not graze or feed livestock on treated areas for six weeks after planting". Add appropriate information for any other chemical treatments.

• • • • • •

.....

.....

•

IMPORTANT NOTICE --- Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions or ust, but neither this warranty nor any other warranty of merchantability or litness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.