


DANGER! KEEP OUT OF REACH OF CHILDREN.

Do not get in eyes.
Causes severe eye damage.
Wear goggles when handling.
In case of contact with eyes, flush immediately with running water and get medical attention.



VITAVAX[®]-EVS CONCENTRATE

ACCEPTED
25 JUL 1975
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 466-124

NET CONTENTS — 5 GALLONS U.S.

COMPOSITION

Active Ingredient: (percent by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide)*	29.52% **
Inert Ingredients:	70.48%
Total	100.00%



* U.S. Patent Nos. 3,249,499; 3,393,202; 3,454,391

** Contains 2.9 pounds of active carboxin per gallon

EPA Reg. No. 400-124 AA

EPA Est. No.

6/11/75

INFORMATION

VITAVAX-EVS concentrate is for use by professional seed treaters only. In this formulation the active ingredient is bound firmly to the seed in a dustfree, water permeable film, that dries quickly without the need of additional heat, improves seed appearance, and eliminates buildup in planter plates. VITAVAX-EVS concentrate is compatible with EVERSIELD SEED PROTECTANT fungicides and insecticides. Observe all cautions and limitations on labeling of all products used in mixtures. Bags containing treated seed should be labeled "Treated seed - do not use for feed, food or oil purposes, or graze or feed livestock on treated areas for six weeks after planting". Add appropriate information for any other chemical treatments.

DIRECTIONS

DO NOT FREEZE.
STORE IN COOL DRY PLACE.

1. Add, at concentrations given in the Table below, Vitavax-EVS Concentrate to cool tap water and mix thoroughly for 15 minutes to prepare the diluted product.
2. Add 7 oz. of EVERSIELD[®] SEED PROTECTANT Liquid dye[†] to each batch of the diluted product and mix for an additional 10 minutes.

Seed	Gallons of Water	Gallons of EVERSIELD Concentrate		Gallons of VITAVAX-EVS Concentrate	Application rates (Cubic centimeters of diluted product per 100 lbs of seed)	Diseases Controlled ³
		Captan ¹	Thiram ²			
Cotton	5	0.7		1	660	<u>Rhizoctonia solani</u>
	6		1	1	785	
	6	0.5	0.5	1	785	
Barley	6			1	335	Loose smut, covered smut
	6		1	1	335	
Oats	4			1	635	Loose smut, covered smut
	4		0.5	0.5	635	
Wheat	6	1		1	325	Loose smut, common bunt
	6		1	1	325	

[†] 21CFR Chapter 1 Section 3.13 requires grain seed treated with poisonous substances in excess of recognized tolerances be suitably colored to prevent their subsequent inadvertent use as food for man or feed for animals.

1. EVERSIELD C SEED PROTECTANT OR EVERSIELD C-M PROTECTANT
2. EVERSIELD T SEED PROTECTANT
3. COMBINATION TREATMENTS PROVIDE CONTROL OF SEED AND SEEDLING DISEASES.

* Evershield is a registered trademark of Cargill, Inc.

DANGER

HANDLING — Harmful if swallowed. Avoid inhaling spray. Wash hands and face thoroughly with soap and water after use and before eating or smoking. Care must be exercised in the handling of treated seed. Keep away from food or food products.

FISH AND WILDLIFE — This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL — Burn or bury container 18 inches below soil surface at safe disposal site. Never reuse for food or water. If disposal is by burning, keep out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, **but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such**

Packaged for Uniroyal Chemical, Division of Uniroyal, Inc., Naugatuck, Connecticut 06770