

470-139

**UNIROYAL**

# UNIROYAL DINOSEB-3

## Dinitro Herbicide

For the control of broadleaf weeds and grasses in many crops

A soluble concentrate containing  
Active ingredient:  
2 sec butyl-4,6 dinitrophenol as the ethanolic salt (of the ethanol and isopropanol series) 51%  
Inert ingredients 49%  
Contains the equivalent of 3 lbs. of 2 sec butyl-4,6 dinitrophenol per gallon

**DANGER - POISON**

**KEEP OUT OF REACH OF CHILDREN**

**MAY BE FATAL IF SWALLOWED OR ABSORBED THROUGH SKIN.**

Do not get in eyes, on skin, or clothing. Avoid breathing spray mist. Do not wear contaminated clothing or shoes. Keep away from farm animals and pets. SEE OTHER WARNINGS AND FIRST AID TREATMENTS ON SIDE PANEL.

Keep away from heat and open flame

**Net Contents 5 Gallons**

Manufactured for  
**UNIROYAL CHEMICAL**  
Division of UNIROYAL INC  
Naugatuck CT 06770

ACCEPTED  
Z...

139

GENERAL INFORMATION

UNIROYAL DINOSEB 3 is a water soluble... herbicide... effective against broadleaf weeds and grasses...

Local conditions affect the use... consult state agricultural experiment stations...

TYPES OF TREATMENT

Preemergence Treatment: Application after the crop is planted but before crop plants emerge... Postemergence Treatment: Application after crop plants emerge...

Preemergence Treatment: Application after crop plants emerge... Postemergence Treatment: Application after crop plants emerge...

CROPPINGS

All applications should be expressed in the basis of water... Soybeans: Preemergence... Postemergence...

Early Postemergence: When soybeans have emerged and are in the early seedling stage... Broadleaf Postemergence Treatment: For this use apply when soybeans are 1 to 2 inches tall...

Broadleaf Postemergence Treatment: For this use apply when soybeans are 1 to 2 inches tall... Note: Do not spray on or over crops or soil...

Note: Do not spray on or over crops or soil...

UNIROYAL DINOSEB 3... Water Soluble Herbicide

PEAS: Preemergence... Postemergence... BASAFACON: Preemergence...

BEANS: Preemergence... Postemergence... CUCURBIT: Preemergence... Postemergence...


LEGUMES: Alfalfa, Lucerne, Clover, Red Clover, Sweet Clover, and Sorghum... CHICKEN: Preemergence... Postemergence...

PEANUTS: Preemergence... Postemergence... Note: Do not graze treated areas before first cutting of hay or within 60 days of treatment.

LYNET: Preemergence... Postemergence... PEANUTS: Preemergence... Postemergence...

PEANUTS: Preemergence... Postemergence... Note: Do not graze treated areas before first cutting of hay or within 60 days of treatment.

400-139



**FRUIT AND NUT ORCHARDS AND GRAPE VINEYARDS IN IRRIGATED AREAS IN CALIFORNIA** - See N.ROYAL DINOSE B 3 at the rate of 1.0 to 2.0 gallons per acre of water per acre to control annual weeds and grasses under conditions specified below. The application rate is based on a 100 gallon of spray per acre. For best results, apply the spray to the weeds and grasses before they are 10 inches high. Later Postemergence - Apply to weeds and grasses after they have emerged from the soil. For best results, apply the spray to the weeds and grasses before they are 10 inches high. For best results, apply the spray to the weeds and grasses before they are 10 inches high.

**POTATOES** - Weed Control - Preemergence - Apply to the soil at the rate of 1.0 to 2.0 gallons per acre of water per acre. For best results, apply the spray to the soil before the potatoes are planted. For best results, apply the spray to the soil before the potatoes are planted.

**SMALL GRAINS** - Barley, Oats, Rye and Wheat - Postemergence - Apply to the weeds and grasses after they have emerged from the soil. For best results, apply the spray to the weeds and grasses before they are 10 inches high.

**STRAWBERRIES** - Apply to the weeds and grasses after they have emerged from the soil. For best results, apply the spray to the weeds and grasses before they are 10 inches high.

**FRUIT AND NUT ORCHARDS AND GRAPE VINEYARDS IN IRRIGATED AREAS IN CALIFORNIA** - See N.ROYAL DINOSE B 3 at the rate of 1.0 to 2.0 gallons per acre of water per acre to control annual weeds and grasses under conditions specified below. The application rate is based on a 100 gallon of spray per acre. For best results, apply the spray to the weeds and grasses before they are 10 inches high. Later Postemergence - Apply to weeds and grasses after they have emerged from the soil. For best results, apply the spray to the weeds and grasses before they are 10 inches high.

**FRUIT AND NUT ORCHARDS AND GRAPE VINEYARDS IN IRRIGATED AREAS IN CALIFORNIA** - See N.ROYAL DINOSE B 3 at the rate of 1.0 to 2.0 gallons per acre of water per acre to control annual weeds and grasses under conditions specified below. The application rate is based on a 100 gallon of spray per acre. For best results, apply the spray to the weeds and grasses before they are 10 inches high. Later Postemergence - Apply to weeds and grasses after they have emerged from the soil. For best results, apply the spray to the weeds and grasses before they are 10 inches high.

480-139

**DAFFODILS, TULIPS, AND SIMILAR FALL PLANTED BULBS:** Pre-emergence use of 1.5 to 2.0 quarts of 4% ROYAL LINDSEY in about 100 gallons of water per acre. Do not apply after bulb shoots emerge above ground.

**GLADIOLUS:** Pre-emergence use of 1.5 to 2.0 quarts of 4% ROYAL LINDSEY in about 100 gallons of water per acre as an overall spray. Preferably just before bulb emergence.

**USE PRECAUTIONS**

To avoid staining carpets, upholstery, etc., keep contaminated clothing and shoes outdoors until cleaned. Rinse equipment and containers with water and dispose of wastes by burying in non-crop lands away from water supplies. Dispose of empty containers by punching holes in them and burying with wastes. This product is toxic to fish and wild life. Keep out of lakes, ponds, and streams. Birds feeding on treated areas may be killed. Do not apply in any area not specified on this label. Do not use in storage areas around the house. Do not use on treated lawns within 4 weeks after treatment.

**POISON**

**SYMPTOMS OF POISONING:** Excessive fatigue, sweating, thirst, and fever. Symptoms of poisoning develop from any type of exposure. **SEE A PHYSICIAN.**

**FIRST AID:** Have patient lie down in a cool place. Give plenty of water.

**IF SWALLOWED:** SEND FOR A PHYSICIAN. Induce vomiting if within 15 minutes. Do not give anything to eat or drink until advised by a physician. Do not give anything to eat or drink until advised by a physician.

**IF SPLASHED IN EYES:** Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention.

**IF SPILLED ON SKIN:** Wash with soap and plenty of water. If on clothing, remove and wash separately. Do not use treated areas until advised by a physician.

**NOTE TO ATTENDING PHYSICIAN:** The active ingredient is 2,4-D. The chemical name is 2,4-Dichlorophenoxyacetic acid.

**WARRANTY AND DISCLAIMER**

This product is sold as a pesticide. It is not a fertilizer. It is not a herbicide. It is not a fungicide. It is not a bactericide. It is not a nematocide. It is not a molluscicide. It is not a rodenticide. It is not an insecticide. It is not a miticide. It is not a acaricide. It is not a nematode. It is not a mollusc. It is not a rodent. It is not an insect. It is not a mite. It is not an acarid. It is not a nematode. It is not a mollusc. It is not a rodent. It is not an insect. It is not a mite. It is not an acarid.

EPA REG. NO. 42513  
EPA EST. NO. 61388-A-01

55