

**PRECAUTIONARY  
STATEMENTS  
HAZARUS TO HUMANS AND  
DOMESTIC ANIMALS  
CAUTION**

AVOID CONTACT WITH SKIN, EYES, OR CLOTHING. WASH HANDS AND FACE THOROUGHLY WITH SOAP AND WATER AFTER USE, AND BEFORE EATING OR SMOKING.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Exposed treated seeds may be hazardous to birds and other wildlife. Dispose of all treated seeds by burial away from bodies of water.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**1. PROHIBITIONS**

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited.

**2. PESTICIDE DISPOSAL**

Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

**3. CONTAINER DISPOSAL**

Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved State and local procedures.

**4. GENERAL**

Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

# VITAVAX-3F

FUNGICIDE

**UNIROYAL**

ACCEPTED

MAR 30 1982

**COMPOSITION**

Active Ingredient (by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathin-3-carboxanilide)*	34.0 **
Inert Ingredients	66.0
Total	100.0

\*U.S. Patents 3,249,499 - 3,393,202 - 3,454,391

\*\*Contains 3.34 pounds of active Carboxin per gallon

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION!**

**STATEMENT OF PRACTICAL TREATMENT**

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled: See a physician if an abnormal reaction occurs.

If on skin: Wash thoroughly with soap and water.

If in eyes: Immediately flush eyes with plenty of water and get medical treatment.

See left side panel for Additional Precautionary Statement.

Mfg. by UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.

Naugatuck, Connecticut 06770

EPA Establishment No.

EPA Registration No. 400-152

NET WEIGHT

**DIRECTIONS FOR USE**

VITAVAX-3F Fungicide is a ready-to-use seed treatment which should be applied as a slurry treatment by machine.

**BARLEY, WHEAT AND OATS**

For the control of loose smut (*Ustilago nuda*), covered smut (*Ustilago hordei*), of barley, loose smut (*Ustilago avenae*), covered smut (*Ustilago kollerii*) of oats, loose smut (*Ustilago tritici*) and common bunt (*Tilletia caries*) of wheat. Use at the rate of 2-3 ounces per 100 pounds of seed.

**CORN**

VITAVAX-3F Fungicide controls seed and seedling diseases and provides systemic control of seedborne head smut (*Sphacelotheca reiliana*). VITAVAX-3F Fungicide is also particularly effective against *Rhizoctonia solani*. VITAVAX-3F Fungicide should be used at a rate of 2 to 4 ounces per 100 pounds of seed when applied to seed that has been treated or will be treated with protectant fungicides such as captan or thiram.

**COTTON**

VITAVAX-3F Fungicide controls seed rots and provides systemic control of seedling diseases of cotton. Seed rot, damping off and sore-shin due to *Rhizoctonia solani* are prevented. VITAVAX-3F Fungicide may be used on both machine and acid delinted cottonseed. Use 3-8 ounces of VITAVAX-3F Fungicide per 100 pounds of seed.

VITAVAX-3F Fungicide can be used on seed previously treated or to be treated with other seed treatment fungicides, such as captan or thiram, or seed treatment insecticides, such as Disyston or Thimet.

**COMPATIBILITY**

VITAVAX-3F Fungicide can be used in combination with Gustafson's CAPTAN 30-DD and CAPTAN-300 or Gustafson's 42-S flowable seed protectants. All other tank mixes should be pre-tested to determine physical compatibility between formulations, and to assure there will be no adverse phytotoxic reactions.

2-17-82

#### FOR CORN SEED ONLY

VITAVAX-3F Fungicide may be tank mixed with the following registered pesticides: Orthocide\* 75 Seed Protectant, Orthocide\* Methoxychlor 75-3 Seed Protectant, Captan 80 Seed Protectant, Captan Methoxychlor 75-2 Seed Protectant, and Cythion\*\* 8-F. Do not allow mixtures to stand overnight. Thoroughly clean equipment between treating operations.

\*Orthocide - trademark of Chevron Chemical Company

\*\*Cythion - trademark of American Cyanamid

Observe all cautions and limitations on labeling of all products used in mixture with this product only as specified on this label.

#### USE RESTRICTIONS

When using formulations that do not contain dye, to comply with 21CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 108.1001) to distinguish and prevent subsequent and inadvertent use as food for man or feed for animals. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Do not graze or feed livestock on treated areas for six weeks after planting.

#### SEED LABELING

"Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Seed commercially treated with VITAVAX-3F Fungicide must be labeled in accord with all applicable requirements of the Federal Seed Act. On corn, barley, and wheat, do not graze or feed livestock on treated areas for six weeks after planting." Add appropriate information for any other chemical treatments.

**IMPORTANT NOTICE** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

1973 Uniroyal, Inc. All Rights Reserved