

PM 13

400-154

1 of 15
file



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

AUG 15 1994

WILLIARD F. CUMMINGS
UNIROYAL CHEMICAL COMPANY, INC.
74 AMITY ROAD
BETHANY, CT. 06524-3402

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 08/18/93 in Response to PR Notice 93-7
EPA Reg. No. 400-154
COMITE II

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,


Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

Willard F. Cummings
Uniroyal Chemical Corporation

Comments for: EPA REG. NO. 400-154
Comite II

The following specific comments pertain to your WPS labeling
submission concerning the product cited above:

Make the changes indicated on the label in red ink.

COMITE® II

AGRICULTURAL MITICIDE

RECOMMENDED FOR AGRICULTURAL USE ONLY

**UNIROYAL
CHEMICAL**

ACCEPTED
with COMMENTS
In EPA Letter Dated

AUG 15 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

400-154

COMPOSITION

Active Ingredients: (% by weight)	
Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]*	68.0%
Inert Ingredients	32.0%
Total:	100.0%

*Contains 6 pounds lbs. technical propargite per g: 'lon

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
EPA REG. NO. 400-154
EPA EST. NO.

005WPS

NET WEIGHT

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.
IF ON SKIN: Remove contaminated clothing and wash skin thoroughly with soap and water. Get medical attention.
IF INHALED: Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.
IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive, causes eye damage. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wash hands and face thoroughly with soap and water after use and before eating drinking or smoking.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; chemical-resistant gloves such as Nitrile or Butyl, barrier laminate, neoprene rubber, polyvinyl chloride or viton; shoes plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing or loading; dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 7 days.

Exception: After the first 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, appears below.

- coveralls
- chemical-resistant headgear
- chemical-resistant gloves, such as nitrile, butyl, barrier laminate, neoprene rubber, polyvinyl chloride or viton
- shoes plus socks
- protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not plant any food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite.

Do not graze or feed livestock on cover crops grown among the trees and vines listed on this label.

Do not mix COMITE II with alkaline materials (such as lime, Bordeaux mixture or lime sulfur) or with materials containing a large amount of petroleum type solvents.

Cotton - Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.

Beans - Do not feed or forage treated bean vines or bean trash after harvesting.

Peanuts - Do not graze or feed livestock on treated areas or cut treated forage for hay.

DIRECTIONS FOR USE

COMITE II is a liquid emulsifiable concentrate for preparing sprays to control the banks grass, Pacific spider, strawberry spider and two spotted spider mites.

COMITE II is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of COMITE II to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

BEST AVAILABLE COPY

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	TIMING OF APPLICATION	AMOUNT OF COMITE II PER ACRE	GALLONS OF SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
				Ground Application	Aerial Application		
COTTON (California & Arizona only)	two-spotted spider Pacific spider strawberry spider	Early ⁽¹⁾	1 1/4 to 1 1/2 pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1 1/4 to 2 1/4 pints	25 ⁽²⁾ to 40	—		
			2 1/4 pints	—	5 to 15		
		Layby	2 1/4 pints	25 ⁽²⁾ to 50	5 to 15		
Layby to Boll Opening	2 1/4 pints	—	5 to 15				
COTTON (East of the Rocky Mts.)	two-spotted spider Pacific spider Strawberry spider	Early ⁽¹⁾	1 1/4 pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1 1/4 to 2 1/4 pints	25 ⁽²⁾ to 40	5 to 15		
		Layby to Boll Opening	2 1/4 pints	25 ⁽²⁾ to 50	5 to 15		
FIELD CORN ^(3,4) (grain & silage)	two-spotted spider Banks grass		2 1/4 pints	minimum of 20	minimum of 2	1	30
GRAIN SORGHUM ⁽⁵⁾ (East of Rocky Mountains)	two-spotted spider Banks grass		1 1/2 to 2 1/4 pints	minimum of 20	minimum of 5	1	30
PEANUTS	two-spotted spider		2 1/4 pints	minimum of 20	minimum of 5	1	14
POTATOES (Pacific Northwest Only)	two-spotted spider		1 1/2 to 2 1/4 pints	20 to 50	minimum of 10	2	14
BEANS ⁽⁶⁾ (green, lima & dry) (West of Rocky Mts.)	two-spotted spider		2 1/4 to 3 1/4 pints	minimum of 20	minimum of 5	2	succulent - 7 dry - 28

APPLICATION NOTES

COTTON

1. COMITE II may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
2. COMITE II may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE II dosage rate per acre.

CORN

3. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. In California, ensure applications are made only when corn leaves are dry. This is especially true for white corn varieties which are the most susceptible to plant injury.
4. In Kansas, Colorado and New Mexico use a minimum of 2 gallons spray solution per acre. For all other states, use a minimum of 5 gallons spray solution per acre.

GRAIN SORGHUM

5. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE II on a large area.

POTATOES

6. The 2 1/2 pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

7. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

® COMITE is a Registered Trademark of Uniroyal Chemical Company, Inc.
© Copyright 1993, Uniroyal Chemical Company, Inc.

BEST AVAILABLE COPY



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

7 of 15

file

AUG 15 1994

WILLIARD F. CUMMINGS
UNIROYAL CHEMICAL COMPANY, INC.
74 AMITY ROAD
BETHANY, CT. 06524-3402

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 08/18/93 in Response to PR Notice 93-7
EPA Reg. No. 400-154
COMITE II

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.

BEST AVAILABLE COPY



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

Willard F. Cummings
Uniroyal Chemical Corporation

Comments for: EPA REG. NO. 400-154
Comite II

The following specific comments pertain to your WPS labeling
submission concerning the product cited above:

Make the changes indicated on the label in red ink.

10 of 15

COMITE® II

AGRICULTURAL MITICIDE

RECOMMENDED FOR AGRICULTURAL USE ONLY



ACCEPTED
with COMMENTS
in EPA Letter Dated

AUG 15 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

400-154

COMPOSITION

Active Ingredients: (% by weight)	
Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]*	68.0%
Inert Ingredients	32.0%
Total:	100.0%

*Contains 6 pounds lbs. technical propargite per gallon

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
EPA REG. NO. 400-154
EPA EST. NO.

005WPS

NET WEIGHT

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.
IF ON SKIN: Remove contaminated clothing and wash skin thoroughly with soap and water. Get medical attention.
IF INHALED: Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.
IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Corrosive, causes eye damage. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wash hands and face thoroughly with soap and water after use and before eating drinking or smoking.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and Other Handlers Must Wear: A long-sleeved shirt & long pants; chemical-resistant gloves such as Nitrile or Butyl, barrier laminate, neoprene rubber, polyvinyl chloride or viton; shoes plus socks; protective eyewear; chemical-resistant headgear for overhead exposure; chemical-resistant apron when cleaning equipment, mixing or loading; dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 7 days.

Exception: After the first 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, appears below.

- coveralls
- chemical-resistant headgear
- chemical-resistant gloves, such as nitrile, butyl, barrier laminate, neoprene rubber, polyvinyl chloride or viton
- shoes plus socks
- protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

USE RESTRICTIONS

Do not apply this product through any type of irrigation system.

Do not plant any food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite.

Do not graze or feed livestock on cover crops grown among the trees and vines listed on this label.

Do not mix COMITE II with alkaline materials (such as lime, Bordeaux mixture or lime sulfur) or with materials containing a large amount of petroleum type solvents.

Cotton - Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.

Beans - Do not feed or forage treated bean vines or bean trash after harvesting.

Peanuts - Do not graze or feed livestock on treated areas or cut treated forage for hay.

DIRECTIONS FOR USE

COMITE II is a liquid emulsifiable concentrate for preparing sprays to control the banks grass, Pacific spider, strawberry spider and two-spotted spider mites.

COMITE II is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control.

After three-quarters full of water, add recommended dosage (from table) of COMITE II to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

BEST AVAILABLE COPY

120715

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	TIMING OF APPLICATION	AMOUNT OF COMITE II PER ACRE	GALLONS OF SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
				Ground Application	Aerial Application		
COTTON (California & Arizona only)	two-spotted spider Pacific spider strawberry spider	Early ⁽¹⁾	1½ to 1¾ pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1½ to 2¼ pints	25 ⁽²⁾ to 40	—		
			2¼ pints	—	5 to 15		
		Layby	2¼ pints	25 ⁽²⁾ to 50	5 to 15		
Layby to Boll Opening	2¼ pints	—	5 to 15				
COTTON (East of the Rocky Mts.)	two-spotted spider Pacific spider Strawberry spider	Early ⁽¹⁾	1¼ pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1½ to 2¼ pints	25 ⁽²⁾ to 40	5 to 15		
		Layby to Boll Opening	2¼ pints	25 ⁽²⁾ to 50	5 to 15		
FIELD CORN ^(3,4) (grain & silage)	two-spotted spider Banks grass		2¼ pints	minimum of 20	minimum of 2	1	30
GRAIN SORGHUM ⁽⁵⁾ (East of Rocky Mountains)	two-spotted spider Banks grass		1½ to 2¼ pints	minimum of 20	minimum of 5	1	30
PEANUTS	two-spotted spider		2¼ pints	minimum of 20	minimum of 5	1	14
POTATOES (Pacific North-west Only)	two-spotted spider		1½ to 2¼ pints	20 to 50	minimum of 10	2	14
BEANS ⁽⁶⁾ (green, lima & dry) (West of Rocky Mts.)	two-spotted spider		2¼ to 3¾ pints	minimum of 20	minimum of 5	2	succulent - 7 dry - 28

APPLICATION NOTES

COTTON

1. COMITE II may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
2. COMITE II may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE II dosage rate per acre.

CORN

3. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. In California, ensure applications are made only when corn leaves are dry. This is especially true for white corn varieties which are the most susceptible to plant injury.
4. In Kansas, Colorado and New Mexico use a minimum of 2 gallons spray solution per acre. For all other states, use a minimum of 5 gallons spray solution per acre.

GRAIN SORGHUM

5. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE II on a large area.

POTATOES

6. The 2¼ pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

7. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.


® COMITE is a Registered Trademark of Uniroyal Chemical Company, Inc.
 © Copyright 1993, Uniroyal Chemical Company, Inc.

BEST AVAILABLE COPY

COMITE® II

AGRICULTURAL MITICIDE

RECOMMENDED FOR AGRICULTURAL USE ONLY



COMPOSITION

Active Ingredients: (% by weight)	
Propargite [2-(p-tert-butylphenoxy) cyclohexyl 2-propynyl sulfite]*	68.0%
Inert Ingredients	32.0%
Total:	100.0%

*Contains 6 pounds technical propargite per gallon

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749
EPA REG. NO. 400-154
EPA EST. NO.

003
10-10-90

NET WEIGHT

KEEP OUT OF REACH OF CHILDREN
DANGER
PELIGRO

NEW STATEMENT

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.
IF ON SKIN: Remove contaminated clothing and wash skin thoroughly with soap and water. Get medical attention.
IF INHALED: Remove person to fresh air. Apply artificial respiration if symptoms indicate. Call a physician.
IF SWALLOWED: Do not induce vomiting. Drink promptly a large quantity of milk, egg whites or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician or Poison Control Center immediately.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS DANGER

Corrosive, causes eye damage. May be fatal if inhaled or absorbed through skin. Harmful if swallowed. Causes skin irritation. Do not breathe vapors or spray mist. Do not get in eyes, on skin or on clothing. Wash hands and face thoroughly with soap and water after use and before eating, or smoking. (See use restrictions for protective clothing and clean-up requirements)

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Flammable, keep away from heat and open flame.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

REENTRY STATEMENT- EXCEPT FOR HARVESTING

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas until sprays have dried. Protective clothing requirements are outlined below for the indicated reentry.

CROP	TIME FOR REENTRY		
	EARLY (1,2) ENTRY WORKER	IRRIGATOR, TRACTOR DRIVER, SCOUT (3)	ALL OTHER (1,4) ACTIVITIES
All labeled crops	when sprays have dried	48 hours	7 days

1. Early entry is defined as any worker who is directed to enter a reentry restricted area prior to the expiration of any reentry interval that

applies to that area. This will include those activities necessitated by unexpected events such as weather, pest conditions, equipment failure, etc. and does not include routine cultural practices.

- These workers will wear protective clothing to include at least:
 - a hat or other suitable head covering
 - long-sleeved shirt and long-legged trousers or a coverall type garment (closely woven fabric covering the body including arms and legs).
 - chemical resistant gloves.
 - shoes and socks.
 NOTE: Occupants of enclosed vehicles must wear normal work attire.
- These workers will wear normal work attire to include at least:
 - a hat or other suitable head covering.
 - long-sleeved shirt and long-legged trousers
 - shoes and socks
- All other activities include those which have high foliar exposure such as pruners, cane turners and tippers (table grapes), etc.
- See use directions for time of harvest after last application. Because certain states may require more restrictive reentry intervals for

various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to the workers who are expected to be in a treated area or in an area to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

"DANGER, Area to be treated with Propargite on (date of application). Do not enter until sprays have dried. Tractor drivers and irrigators and scouts wearing normal work attire may enter after 48 hours. Do not enter treated areas without protective clothing to perform activities that involve substantial or prolonged exposure with treated foliage for (enter number of days)."

If accidentally exposed, consult the Statement of Practical Treatment portion of this label for first aid recommendations.

USE RESTRICTIONS

Mixer/loaders must wear goggles or a face shield, chemical-resistant apron, long-sleeved shirt, long pants, and mid-forearm to elbow length chemical-resistant gloves.

Any article of clothing worn while mixing, loading or applying product must be cleaned before re-use. Clothing should be laundered separately from household articles. Clothing that has been drenched or has otherwise absorbed concentrated pesticide must be disposed of in a sanitary landfill, incinerated, or burned if allowed by State and local authorities. Do not apply this product through any type of irrigation system.

Do not plant any food or feed crop in rotation within 6 months after last application of propargite unless the crop is a registered use for propargite. Do not graze or feed livestock on cover crops grown among the trees and vines listed on this label.

Do not mix COMITE II with alkaline materials (such as lime, Bordeaux mixture or lime sulfur) or with materials containing a large amount of petroleum type solvents.

Cotton - Apply only before bolls open. Do not feed treated foliage or cotton trash to livestock.

Beans - Do not feed or forage treated bean vines or bean trash after harvesting.

Peanuts - Do not graze or feed livestock on treated areas or cut treated forage for hay.

DIRECTIONS FOR USE

COMITE II is a liquid emulsifiable concentrate for preparing sprays to control the banks grass, citrus red, citrus rust, false spider, Pacific spider, six-spotted, strawberry spider, Texas citrus, and two-spotted spider mites.

COMITE II is not systemic in action, therefore complete coverage of both upper and lower leaf surfaces and of fruit is necessary for effective control. After three-quarters full of water, add recommended dosage (from table) of COMITE II to the spray tank. Fill tank, agitate and spray thoroughly to cover foliage and fruit for best results.

BEST AVAILABLE COPY

DOSAGE INSTRUCTIONS

CROP	MITES CONTROLLED	TIMING OF APPLICATION	AMOUNT OF COMITE II PER ACRE	GALLONS OF SPRAY SOLUTION PER ACRE		TOTAL NUMBER OF SPRAYS PER YEAR	DAYS BEFORE HARVEST (LAST SPRAY)
				Ground Application	Aerial Application		
COTTON (California & Arizona only)	two-spotted spider Pacific spider strawberry spider	Early ⁽¹⁾	1 1/4 to 1 1/2 pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1 1/4 to 2 1/4 pints	25 ⁽²⁾ to 40	—		
			2 1/4 pints	—	5 to 15		
		Layby	2 1/4 pints	25 ⁽²⁾ to 50	5 to 15		
Layby to Boll Opening	2 1/4 pints	—	5 to 15				
COTTON (East of the Rocky Mts.)	two-spotted spider Pacific spider Strawberry spider	Early ⁽¹⁾	1 1/4 pints	15 ⁽²⁾ to 30	Not recommended	3	Before bolls open
		Mid Season to Layby	1 1/4 to 2 1/4 pints	25 ⁽²⁾ to 40	5 to 15		
			Layby to Boll Opening	2 1/4 pints	25 ⁽²⁾ to 50		
FIELD CORN ⁽³⁾ (grain & silage)	two-spotted spider Banks grass		2 1/4 pints	minimum of 20	minimum of 2	1	30
GRAIN SORGHUM ⁽³⁾ (East of Rocky Mountains)	two-spotted spider Banks grass		1 1/4 to 2 1/4 pints	minimum of 20	minimum of 5	1	30
PEANUTS	two-spotted spider		2 1/4 pints	minimum of 20	minimum of 5	1	14
POTATOES (Pacific Northwest Only)	two-spotted spider		1 1/4 to 2 1/4 pints	20 to 50	minimum of 10	2	14
BEANS ⁽⁴⁾ (green, lima & dry) (West of Rocky Mts.)	two-spotted spider		2 1/4 to 3 1/4 pints	minimum of 20	minimum of 5	2	succulent - 7 dry - 28

APPLICATION NOTES

COTTON

- 1. COMITE II may be phytotoxic to young cotton under certain conditions. Age and climatic conditions result in a physiological hardening when plants are generally between 6 and 12 inches tall. Applications can begin after plants are hardened.
- 2. COMITE II may be used in concentrate sprayers designed for concentrate row crop applications. Maintain COMITE II dosage rate per acre.

CORN

- 3. Treat early. Best results are achieved when mite populations are low, before the mites have damaged the plants. Treatment is recommended when mite populations are just beginning to build, and the corn is still small enough (usually 3 to 4 feet) to allow good coverage. In California, ensure applications are made only when corn leaves are dry. This is especially true for white corn varieties which are the most susceptible to plant injury.
- 4. In Kansas, Colorado and New Mexico use a minimum of 2 gallons spray solution per acre. For all other states, use a minimum of 5 gallons spray solution per acre.

GRAIN SORGHUM

- 5. Due to the large number of varieties of sorghum, it is recommended that a small test plot be treated to check phytotoxicity before using COMITE II on a large area.

POTATOES

- 6. The 2½ pints per acre rate should be used when the crop is under extreme mite pressure.

BEANS

- 7. Do not use on varieties grown for fresh market or on varieties whose value is linked to pod appearance as pods may be spotted.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

IMPORTANT NOTICE—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

® COMITE is a Registered Trademark of Uniroyal Chemical Company, Inc.

© Copyright 1984 Uniroyal Chemical Company, Inc.

BEST AVAILABLE COPY