

VITAVAX[®]-M

FLOWABLE FUNGICIDE
(VITAVAX AND THIRAM PLUS MOLYBDENUM)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION

Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid inhaling of spray mist. Do not use or store near food or feed.

During commercial exposure in seed treatment and loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety and Health Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under provisions of 30 CFR, Part 11. Wash hands, arms and face thoroughly with soap and water after handling and before eating or smoking. Routinely shower or bathe after work; wash all clothing with soap and water before reusing. Consumption of alcoholic beverages increases the toxic effects of thiram.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

UNIROYAL

ACCEPTED

JUN 27 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
approved for the pesticide
EPA Reg. No. 400-156

COMPOSITION

Active Ingredients: (% by weight)	
Carboxin (5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide)*	5.7%
Thiram (tetramethylthiuram disulfide)	5.7%
Inert Ingredients:	88.6%
Total:	100.0%

*U.S. Patents 3,249,499 - 3,393,202 - 3,454,391

**Contains 2.90% of elemental Molybdenum derived from sodium molybdate.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and see a physician.

IF INHALED: See a physician if an abnormal reaction occurs.

IF ON SKIN: Wash thoroughly with soap and water.

IF IN EYES: Immediately flush eyes with plenty of water. See a physician.

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

CHEMICAL GROUP — Uniroyal, Inc.

Middlebury, CT 06749

EPA REG. NO. 400-156

EPA EST. NO.

VITAVAX is a Registered Trademark of UNIROYAL, Inc.

Copyright 1985, Uniroyal, Inc.

NET WEIGHT:

8-17-85

Int. No.

DIRECTIONS FOR USE

VITAVAX-M Flowable Fungicide is a ready-to-use seed treatment. VITAVAX-M Flowable Fungicide combines the systemic action of carboxin with the surface action of thiram to control various seed and seedling diseases of soybeans. It is particularly effective against *Rhizoctonia solani*. VITAVAX-M Flowable Fungicide also contains the micronutrient molybdenum.

VITAVAX-M Flowable Fungicide can be applied as a Hopper Box treatment or in a mechanical seed treater.

SOYBEANS

Use 6 fl. oz. of VITAVAX-M Flowable Fungicide per bushel of seed or 12 fl. oz. per 100 pounds of seed. Six (6) to twelve (12) fl. oz. of VITAVAX-M Flowable Fungicide will produce the equivalent of 0.2 - 0.4 oz. of elemental molybdenum.

Mix well before use. Apply VITAVAX-M Flowable Fungicide directly to seed in hopper box. Partially fill hopper box with a premeasured amount of seed. Apply the proper amount of VITAVAX-M Flowable Fungicide evenly over the surface of the seed. Mix with a tick or paddle until a uniform color is imparted to all seed. Repeat procedure until hopper box is filled.

For mechanical seed treaters apply VITAVAX-M Flowable Fungicide at a dosage of twelve fl. oz. per 100 pounds of seed directly to a premeasured amount of seed and mix thoroughly until all seed have a uniform color.

USE RESTRICTIONS

Do not graze or feed livestock on vines grown from treated seed. Care must be exercised in the handling of treated seed.

USE PRECAUTION

When using formulations that do not contain dye, to comply with 21 CFR Chapter 1, Section 2.25, all grain seed treated with an economic poison must be colored with an approved dye (40 CFR 180.1001) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals.

COMPATIBILITY

VITAVAX-M Flowable Fungicide should not be tank mixed with any pesticide unless adequate physical compatibility and phytotoxicity tests have been completed and proved to be satisfactory.

SEED LABELING

"This seed treated with VITAVAX-M Flowable Fungicide. Do not use treated seed for food, feed, or oil purposes. Add appropriate information for any other chemical treatments."

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

