

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS  
CAUTION**

Avoid inhaling dust. Wear respirator. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. WASH HANDS AND FACE THOROUGHLY WITH SOAP AND WATER AFTER USE, AND BEFORE EATING OR SMOKING.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

This is a violation of Federal law to use this product in a manner inconsistent with the labeling.

**STORAGE AND DISPOSAL**

- 1. PROHIBITIONS**  
Do not contaminate water, food, or feed. Do not spray directly on food. Do not use equipment used for spraying in other ways.
- 2. PESTICIDE DISPOSAL**  
Pesticide spray mixture and residue that is left in the container should be disposed of in accordance with the Federal State and local regulations. For more information, contact the Resource Conservation and Recovery Act.
- 3. CONTAINER DISPOSAL**  
**Bags:** Do not burn. Contact your local health department for disposal instructions.  
**Fiber Drum:** Do not burn. Contact your local health department for disposal instructions.
- 4. GENERAL**  
Observe Federal, State, and local regulations for disposal of this product.
- 5. STORAGE**  
Store in original container.

# UNIROYAL Captan 50W

FUNGICIDE



**COMPOSITION**

<b>ACTIVE INGREDIENT</b>	
Captan 50W (50% wettable powder)	4.00
<b>INERT INGREDIENTS</b>	96.00
<b>Total</b>	100.00

**DIRECTIONS FOR USE**

Apply the recommended amount of this material on surface of water in nearly full spray tank or pre-mix in a bucket. Mix with water and pour mixture through screen into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to starve.

Consult state Agricultural Extension Service authorities for specific information as the timing, number and rate of applications will vary with local conditions.

**KEEP OUT OF REACH OF CHILDREN.  
CAUTION**

**STATEMENTS OF PRACTICAL TREATMENT**

If swallowed: Induce vomiting and see a physician.  
If inhaled: See a physician if an abnormal reaction occurs.  
If on skin: Wash thoroughly with soap and water.  
If in eyes: Immediately flush eyes with plenty of water. See a physician.  
See left side panel for additional precautionary statement.

Mfg. by UNIROYAL CHEMICAL, Division of UNIROYAL INC.  
Naugatuck, Connecticut 06770  
EPA Establishment No. 904 NJ 2  
EPA Registration No. 400 157

NET WEIGHT

# UNIROYAL Captan 50W

FUNGICIDE



COMPOSITION

ACTIVE INGREDIENT	
Captan (N-Trichloromethyl-4-chlorobenzamide)	50.0
INERT INGREDIENTS	50.0
Total	100.0

DIRECTIONS FOR USE

Pour recommended amount of material on surface of water in nearly filled spray tank or premix in a bucket half filled with water and pour mixture through screen into nearly filled spray tank. Add balance of water to fill tank. Keep agitating during filling and spraying operations. Do not allow mixture to stand.

Consult state Agricultural Extension Service authorities for specific information as to timing, number and rate of applications which vary with local conditions.

KEEP OUT OF REACH OF CHILDREN.

**CAUTION**

STATEMENTS OF PRACTICAL TREATMENT

- If swallowed: Induce vomiting and see a physician
- If inhaled: See a physician if an abnormal reaction occurs
- If on skin: Wash thoroughly with soap and water
- If in eyes: Immediately flush eyes with plenty of water. See a physician
- See left side panel for additional precautionary statement

Mfg. by UNIROYAL CHEMICAL, Division of UNIROYAL, INC.  
Naugatuck, Connecticut 06770  
EPA Establishment No. 904 NJ 2  
EPA Registration No. 400 157

NET WEIGHT

1 of 2

**PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

Avoid inhaling dust. Wear respirator. AVOID CONTACT WITH SKIN, EYES OR CLOTHING. WASH HANDS AND FACE THOROUGHLY WITH SOAP AND WATER AFTER USE, AND BEFORE EATING OR SMOKING.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply to water. Do not contaminate water by use of equipment or disposal of waste.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

See left side panel for directions for use.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. For cleaning purposes, use only open dumping is prohibited.

PESTICIDE DISPOSAL

Do not spray mixture of this material. Do not be used according to label instructions. Must be disposed according to Federal, State, or local laws under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Do not reuse containers. Do not reuse containers. Do not reuse containers. Do not reuse containers.

OPERATOR DRUGS

Do not use this material if you are under the influence of any drug, alcohol, or other substance which may impair your ability to use this material safely.

GENERAL

Read and follow the label instructions. Do not use this material if the label is missing or illegible.

STORAGE

Store in a cool, dry place.



### COMPATIBILITY

50-W can be used at recommended rates daily with spray schedules in recommended Experiment Station papers. Do not combine and strongly alkaline materials such as hydrated lime which reduces the fungicidal activity. Do not use product in combination with insecticides, particularly following an insect spray, or with soil treatments of Parathion products.

tan 50-W is a microfine wettable powder for use as a spray for the control of certain fungi on fruit, vegetables and ornamental crops and as a fungicide for the control of certain seed rot and damping-off diseases. Spray injury may result if applied immediately before or after a wettable powder spray.

### RECOMMENDATIONS

Usage rates on this label indicate the amount of product per 100 gallons of dilute spray solution.

**APPLE** — scab. For primary infection, apply 1 to 2 lbs. in prebloom, blossom, leaf fall and harvest periods. For secondary infection, apply 1 to 2 lbs. Apply in cover periods. **BERRY** — *Botrytis rot*, *leaf spot*, *black rot*, *botrytis rot*. 1 to 2 lbs. per 100 gallons. Apply in late cover and prebloom periods. **PEACH** — *Botrytis rot*, *blight*. 2 to 3 lbs. per 100 gallons. Apply in first and second cover sprays and prebloom, harvest sprays or as needed at 5 to 7 day intervals. **PUMPKIN** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom through early cover periods. **SPINACH** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. **STRAWBERRY** — *Botrytis rot*. 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. **WATERMELON** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. **WINTER SQUASH** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. **ZUCCHINI** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. **Other uses:** Apply 1 to 2 lbs. per 100 gallons in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**BERRY** — *Botrytis rot*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**PEACH** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**PUMPKIN** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**PEAR** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**PLUMS PRUNES** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**RASPBERRIES** — *Botrytis rot*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**STRAWBERRIES** — *Botrytis rot*. 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

### VEGETABLE CROP USES

**ASPARAGUS** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**CANTALOUPE CUCUMBER PUMPKIN SUMMER and WINTER SQUASH WATERMELON** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**POTATOES** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**POTATO SEED PIECE TREATMENT** — 1 to 2 lbs. per 100 gallons. Apply to seed pieces before planting. Repeat applications at 7 to 14 day intervals.

**POTATO STORAGE ROT CONTROL** — 1 to 2 lbs. per 100 gallons. Apply to potato tubers before storage. Repeat applications at 7 to 14 day intervals.

**TOMATO** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

### OTHER USES

**PACKING BOXES** — 1 to 2 lbs. per 100 gallons. Apply to packing boxes before use. Repeat applications at 7 to 14 day intervals.

**SOIL AND GREENHOUSE BENCH (REPLANTING TREATMENT)** — 1 to 2 lbs. per 100 gallons. Apply to soil and benches before planting. Repeat applications at 7 to 14 day intervals.

**AZALEAS CARNATIONS CHRYSANTHEMUM** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**CARNATION** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**GRASSES** — *downy mildew*. 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

### SEED TREATMENT

**BEANS CABBAGE CORN MELONS PEAS SQUASH** — 1 to 2 lbs. per 100 gallons. Apply to seeds before planting. Repeat applications at 7 to 14 day intervals.

**BEETS CHARD SPINACH GRASS** — 1 to 2 lbs. per 100 gallons. Apply in prebloom and early cover periods. Repeat applications at 7 to 14 day intervals.

**IMPORTANT NOTICE:** The manufacturer warrants that this product is suitable for the uses specified on the label. The manufacturer does not warrant the merchantability or fitness for any particular purpose, express or implied, or extends to the use of this product contrary to the instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.