

UNIROYAL DINOSEB-3.3

A CONTACT WEED KILLER
WITH ADDED SURFACTANT — CONTAINS NITRATED PHENOLS

UNIROYAL

COMPOSITION

Active Ingredients: (% by weight)	
dinoseb (2-sec-butyl-4,6-dinitrophenol)	38.3%
Inert Ingredients:	61.7%
Total:	100.0%

Contains the equivalent of 3.3 lbs. of DINOSEB per gallon.

DIRECTIONS FOR USE

UNIROYAL DINOSEB 3.3 when used with the proper amount of water, will control most of the common herbaceous weeds and grasses. Use enough spray to wet all weed growth thoroughly, as it acts by contact action only. The lower dosages will be satisfactory for the control of small seeding annual grasses such as crabgrass, foxtail, and barnyard (water) grass, and of small seeding annual weeds such as pigweed, mustard, lambsquarter, and sorrel. Coarse rayweed mallow, pineapple weed, sweet fennel and wild carrot will require the higher dosages, particularly when mature and hard to kill.

NOTE: Contact weed killers kill annuals, but perennials must be retreated as new growth develops. Control will be best in warm weather.

MIXING INSTRUCTIONS: Fill sprayer tank $\frac{3}{4}$ full. Add required amount of UNIROYAL DINOSEB 3.3, then add the remaining required amount of water. Agitation must be maintained during mixing and application to keep the spray uniformly mixed.

GENERAL WEED CONTROL: For use in non-crop areas such as industrial sites, around buildings, along ditch banks, and utility, highway, pipeline and railroad rights-of-way, use 3 to 4 $\frac{1}{2}$ pints of UNIROYAL DINOSEB 3.3 per 100 gallons of water. Spray weed growth thoroughly.

ORCHARD FLOORS: ALMONDS, APPLES, APRICOTS, CHERRIES, CITRUS, DATES, FILBERTS, PEACHES, PEARS, PECANS, PLUMS, PRUNES, WALNUTS: Use 3 to 4 $\frac{1}{2}$ pints UNIROYAL DINOSEB 3.3 in 100 gallons of water. Apply as a direct spray to ground cover. Completely cover all weed foliage. Do not apply more than 100 gallons of spray per acre. When applying to orchard floors avoid getting any of the spray on fruits or foliage. Destroy any fruit accidentally sprayed. Avoid spraying the base of the young trees as girdling may occur. Do not allow livestock to graze on treated ground cover.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, wildlife and birds. Birds and other wildlife in treated areas may be harmed. Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act. **CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

PHYSICAL OR CHEMICAL HAZARDS

FLAMMABLE LIQUID: Keep away from Heat and Open Flame. Protect from temperature below 15 degrees F.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

KEEP OUT OF REACH OF CHILDREN DANGER

STATEMENT OF PRACTICAL TREATMENT

INTERNAL: To induce vomiting give a tablespoon of baking soda dissolved in a glass of warm water. Repeat until vomit fluid is clear. CALL A PHYSICIAN IMMEDIATELY.

EXTERNAL: Wash with rubbing alcohol and follow with a thorough scrubbing with soap and water. Remove contaminated clothing immediately. If splashed in eyes, immediately flush with plenty of water for at least 15 minutes and get medical attention.



POISON



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

POISONOUS IF SWALLOWED OR ABSORBED THROUGH SKIN. DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. In case of contact, remove clothing and wash thoroughly with soap and water. FOR EYES: Flush with water and get medical attention. KEEP AWAY FROM CHILDREN, DOMESTIC ANIMALS and FOODSTUFFS.

NOTE TO PHYSICIAN: Active ingredient is a metabolic stimulant. Treat symptomatically. Not for Use or Storage in or Around the Home

See side panel for additional precautionary statements

UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.
Nauvau, Cor. Icut 08770
EPA REG. NO. 4. 158
EPA EST. NO.

NET CONTENTS:

3-18-82

WEED CONTROL IN GRAPES AND BUSH FRUITS: Apply spray containing UNIROYAL DINOSEB 3.3 when weeds are growing well but before they are 6 inches high. For use on grapes in the spring before bloom or in the fall after harvest use 3-4 pints of UNIROYAL DINOSEB 3.3 in 100 gals. of water. For treating during bloom and up to 3 to 4 weeks after bloom use no more than 3 pints of UNIROYAL DINOSEB 3.3 per 100 gallons of water. Use up to 150 gallons per acre to give good spray coverage. Apply only as a directed spray to the weeds and soil in such a way as to avoid contacting grape foliage, blooms, or fruiting clusters. It is often desirable to limit spray application to a 1-2 foot band directly under the grape trellis where weed control problem is most severe and soil cannot be easily cultivated. For such band treatments the gallonage per acre should be reduced proportionally. Make one or two applications as needed.

On Bush Fruits such as blackberries, blueberries, currants, gooseberries, and raspberries, use 3 to 4½ pints of UNIROYAL DINOSEB 3.3 per 100 gallons of water. Apply at the rate of 125 to 150 gallons of spray per acre in the fall after harvest or in the spring before bloom. Use directed spray and do not treat young canes desired for permanent plants. Make one or two repeat applications as needed.

WEED CONTROL IN ALFALFA: Established stands, dormant season; (for control of winter annuals). Use 3 pints UNIROYAL DINOSEB 3.3 in 100 gallons of water to insure adequate coverage. **DO NOT GRAZE TREATED AREAS OR USE TREATED FOLIAGE FOR FEED OR FORAGE.** Do not exceed 4½ pints per acre.

POTATO VINE KILLING: (Seed and table stock) **HIGH VOLUME SPRAYS:** Use 3 pints of UNIROYAL DINOSEB 3.3 in water to make 100 gallons. Use 100 gallons of spray per acre. **MEDIUM VOLUME SPRAYERS:** Use 3 pints UNIROYAL DINOSEB 3.3 in water to make 25 to 40 gallons of water

per acre. **LOW VOLUME SPRAYERS:** Use 3 pints of UNIROYAL DINOSEB 3.3 in 20 gallons of water per acre, apply by either air or ground equipment. Fly a maximum swath of 15 feet. Apply 10 to 20 days before harvest, spraying thoroughly as complete coverage is essential. Do not spray within 10 days of harvest. Do not spray exposed tubers. Do not graze treated areas.

NOTE: Browning of the vascular ring of potato tubers sometimes occurs after the use of vine killers.

Do not allow the material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant, or crop injury, illegal residues or other undesirable results may occur.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.