

PR 93 400-165 138

*File*



**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**  
WASHINGTON, D.C. 20460

**MAR. 30 1994**

**OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES**

Allen R. Blem  
UNIROYAL CHEMICAL CO. INC.  
74 AMITY RD  
BETHANY, CT 06524

**Subject: Label Amendment Submission of 08/09/93 in Response to PR Notice 93-7  
EPA Reg. No. 400-165  
ROYAL MH-30 SG**

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

**WHAT THIS ACCEPTANCE MEANS:**

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

**WHAT YOU NEED TO DO NEXT:**

Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



**Recycled/Recyclable**  
Printed with Soy/Canola Ink on paper that  
contains at least 50% recycled fiber

258  
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division

Allen R Blem  
UNIROYAL CHEMICAL CO INC  
74 AMITY RD  
BETHANY CT 06524

Comment for: EPA Reg Nr.400-165  
ROYAL MH-30 SG

The following specific comments pertain to your WPS  
labeling submission concerning the product  
cited above:

No comment.

# ROYAL MH-30 SG®

## WATER SOLUBLE GRANULE GROWTH RETARDANT FOR POTATOES AND ONIONS

**UNIROYAL  
CHEMICAL**

### COMPOSITION

Active Ingredient: (% by weight)	
Potassium salt of 1,2-dihydro-3,6-pyridazinedione .....	80.0%**
Inert Ingredients .....	20.0%
Total: .....	100.0%

\*\*Contains 60% by weight of maleic hydrazide

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749  
EPA REG. NO. 400-165  
EPA EST. NO.

003WPS

### NET CONTENTS

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Flush eyes with plenty of water. Call a physician if irritation persists.

**IF ON SKIN:** Wash with plenty of soap and water.

**IF INHALED:** Remove to fresh air.

### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury. Avoid contact with skin, eyes or clothing.

### PERSONAL PROTECTIVE EQUIPMENT

**Applicators and Other Handlers Must Wear:** Long-sleeved shirt and long pants; waterproof gloves; shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

coveralls; waterproof gloves; shoes plus socks

**BEST AVAILABLE COPY**

## NOTES

1. Do not apply this product through any type of irrigation system.
2. Follow dosage recommendations carefully and spray field uniformly.
3. Do not use on potato plants grown for seed production.
4. Even though ROYAL MH-30SG is non-volatile no drift should be permitted onto nearby crops. Particular care should be taken to prevent drift when aerial applications are being made.
5. Spray potatoes or onions at least 24 hours before rain or applying overhead irrigation. Rain or overhead irrigation occurring within 24 hours after application will reduce effectiveness.
6. At least a 2-week interval should be allowed between a ROYAL MH-30SG application to potatoes and the use of a vine killer.
7. Uneven application may result in budend cracking and/or elephant hide on potato tubers.
8. To obtain desired results plants should not be stressed from either too little or excessive water before or after application.
9. **IMPORTANT: Geographical response of potato plants.**

### EAST OF THE ROCKY MOUNTAINS

Occasional injury to the terminal growth of potato foliage has been observed following the application of ROYAL MH-30SG. However, trials and usage indicate that this does not adversely affect the yield or the quality of the potato tubers at harvest if the chemical is applied according to label recommendations.

### WEST OF THE ROCKY MOUNTAINS

Reduced sizing, yield, tuber cracking and /or "elephant hide" disorders have been observed in some instances following application of ROYAL MH-30SG on potatoes grown under irrigation in the Northwest. At this time, the association, if any, between these disorders and ROYAL MH-30SG use is not clear or confirmed. Until such time as the association or the lack of association can be determined, ROYAL MH-30SG is not recommended for use on irrigated potatoes, and use on any potatoes, irrigated or not, is at the user's risk. Uniroyal will not be responsible for any loss incurred by users should any of these disorders occur after the application of ROYAL MH-30SG.

10. Do not add any extra wetting agents or commercial spray adjuvants to ROYAL MH-30SG spray solutions except as specified under potatoes and onions.

### MIXING INSTRUCTIONS

**NOTE:** ROYAL MH-30SG must be dissolved first if it is to be tank mixed with an insecticide or fungicide.

1. Be sure sprayer is clean and not contaminated with any material.
2. Fill tank  $\frac{3}{4}$  full with clean water, and start agitation.
3. Pour ROYAL MH-30SG slowly from bag into tank. Avoid dumping contents all at once.
4. Allow 2-3 minutes agitation for dissolving.
5. Add compatible tank mix insecticide or fungicide if needed.
6. Complete filling tank.

### DIRECTIONS FOR USE

#### ONIONS

#### TO CONTROL SPROUTING OF ONIONS IN STORAGE

##### Timing

ROYAL MH-30SG should be applied when the bulbs are fully mature, have at least 5 to 8 green leaves, and necks soft enough for the tops to fall if they have not already done so. The spray is often applied when 50 percent of the tops have fallen but all tops are still green. This condition usually occurs about two weeks to 10 days before harvest time. Green tops are essential for the absorption of ROYAL MH-30SG.

##### Rate of Application

**NOTE:** Uniform and complete coverage is essential for good sprout control.

##### Ground Application

Use  $3\frac{1}{2}$  pounds of ROYAL MH-30SG in a minimum of 30 gallons of water per acre.

##### Aerial Application

Use  $3\frac{1}{2}$  pounds of ROYAL MH-30SG in a minimum of 10 gallons of water per acre. Do not fly at heights of less than 8-12 feet. Maintaining this height will give more uniform coverage and aid in preventing an excessive build-up of chemical on the left side of the treated swath. These heights should be reduced as cross-wind velocities exceed 2-4 MPH.

##### Spray Adjuvants

**West of Rocky Mountains:** Use Regulad, X-77 or R11 at the rate of 1 quart per 100 gallons of spray solution in arid sections of the U.S. These products will help provide better coverage and aid in translocation of the ROYAL MH-30SG.

**East of Rocky Mountains:** use of surfactants is normally not recommended.

##### Temperature

ROYAL MH-30SG should be applied at temperatures below 85°F or at temperatures below 80°F if the temperature is expected to exceed 85°F. ROYAL MH-30SG is water soluble and under high temperature and low humidity conditions will crystallize on the leaf surface and not be absorbed into the plant.

##### CAUTION

Avoid spraying edible onions too early. Spongy bulbs may result if spraying is done earlier than two weeks before maturity.

##### Varieties

Use of ROYAL MH-30SG should be combined with selection of a good-keeping variety. Do not apply to onion varieties with poor-keeping quality. ROYAL MH-30SG will not make poor-keeping varieties equal to good-keeping varieties.

## POTATOES

### TO IMPROVE QUALITY AND TO CONTROL SPROUTING IN STORAGE

#### BENEFITS

##### Sprout Control

ROYAL MH-30SG will control sprouting of potatoes in storage. However, ROYAL MH-30SG will not make a poor-keeping potato such as Norgold or Norchip into a good-keeping potato.

##### Reduction of Volunteer Potatoes

ROYAL MH-30SG will reduce the number of volunteer potatoes the following spring because of its ability to control sprouting. If applied at the proper time it will also reduce the number of undersize potatoes left in the field, which also reduces the potential for volunteers.

##### May Improve Grade

ROYAL MH-30SG may improve the grade of potatoes, especially Russet Burbank variety, if the application is made prior to any secondary tuber formation.

##### Undersize Potatoes

ROYAL MH-30SG will help reduce any additional "sets" of potatoes, thus decreasing the number of undersize potatoes in the field at harvest.

##### Reduces Shrinkage

ROYAL MH-30SG applications can reduce shrinkage in storage up to 15 percent.

### DIRECTIONS FOR USE

##### Timing

Timing is important. To allow translocation of chemical down to the tubers ROYAL MH-30SG should be applied after tubers are formed and while vines are still green and are free from insects, diseases or other damage. Time of application may occur as early as 2-3 weeks past full bloom when the primary and secondary inflorescences have lost most of their flowers and a few third order inflorescences and buds are

**BEST AVAILABLE COPY**

present. This occurs when a few lower leaves show a yellow color. Application prior to this time may result in reduced size of potatoes and phytotoxicity. The following tuber sizes may also be used as a guide, provided the above conditions have been met.

#### **Russet Burbank**

The smallest tubers that are required to reach a minimum size of 4 - 6 oz. must be 1½ - 2" in diameter (and elongating) at time of spraying.

#### **Norgold or Norchip**

The smallest tubers that are required to reach maximum size must be 2" or more in diameter at time of spraying.

#### **Other Varieties**

The smallest tubers that are required to reach maximum size, should be 1½" in diameter at time of spraying.

#### **Rate of Application**

NOTE: Uniform and complete coverage is essential for satisfactory results.

#### **Ground**

Use 5 pounds of ROYAL MH-30SG in a minimum of 30 gallons of water per acre.

#### **Aerial**

Use 5 pounds of ROYAL MH-30SG per acre.

West of the Rocky Mountains: Use in a minimum of 10 gallons of water per acre.

East of the Rocky Mountains: Use in a sufficient quantity of water to gain complete uniform coverage.

Do not fly at heights of less than 8-12 feet. Maintaining this height will give more uniform coverage and aid in preventing an excessive build up of chemical on the left side of the treated swath. These heights should be reduced as cross-wind velocities exceed 2 - 4 MPH.

#### **Spray Adjuvants**

West of Rocky Mountains: Use Regulaid, X-77 or R11 at the rate of 1 quart per 100 gallons of spray solution in arid sections of the U.S. These products will help provide better coverage and aid in translocation of the ROYAL MH-30SG.

East of the Rocky Mountains: Use of surfactants is normally not recommended.

#### **Temperature**

ROYAL MH-30SG should be applied at temperatures below 85°F or at temperatures below 80°F if the temperature is expected to exceed 85°F. later that day. ROYAL MH-30SG is water soluble and under high temperature and low humidity conditions will crystallize on the leaf surface and not be absorbed into the plant.

## **STORAGE AND DISPOSAL**

**Do not contaminate water, feed or feed by storage or disposal.**

**STORAGE:** Store in a dry location.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinseate is a violation of Federal Law. If these wastes can not be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**EQUIPMENT CLEANING:** ROYAL MH-30SG is not corrosive to spray equipment. Following applications, all spray equipment should be thoroughly rinsed with water.

**CONTAINER DISPOSAL:**

**Fiber carton with liners or bags:** Completely empty liner or bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of in a sanitary landfill, or by incineration if permitted by State and local authorities. Dispose of carton in the same manner.

**IMPORTANT NOTICE**—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

© ROYAL MH-30SG is a Registered Trademark of Uniroyal Chemical Company, Inc.

© Copyright 1993, Uniroyal Chemical Company, Inc.

**BEST AVAILABLE COPY**

# ROYAL MH-30 SG<sup>®</sup>

WATER SOLUBLE GRANULE

GROWTH RETARDANT

FOR TOBACCO

**UNIROYAL  
CHEMICAL**

COMPOSITION

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

MAR 30 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

400-165

Active Ingredient: (% by weight)	
Potassium salt of 1,2-dihydro-3,6-pyridazinedione .....	80.0%**
Inert Ingredients .....	20.0%
Total: .....	100.0%

\*\*Contains 60% by weight of maleic hydrazide

UNIROYAL CHEMICAL COMPANY, INC., Middlebury, CT 06749  
EPA REG. NO. 400-165  
EPA EST. NO.

003WPS

NET CONTENTS

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.  
IF ON SKIN: Wash with plenty of soap and water.  
IF INHALED: Remove to fresh air.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury. Avoid contact with skin, eyes or clothing.

#### PERSONAL PROTECTIVE EQUIPMENT

Applicators and Other Handlers Must Wear: Long-sleeved shirt and long pants; waterproof gloves; shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks

**BEST AVAILABLE COPY**

## NOTES

1. To control tobacco suckers apply ROYAL MH-30SG at the proper time. Too early spraying may prevent normal development of top leaves. Too late spraying will allow some sucker growth. FULL FLOWER is the stage when 90% of your plants have begun to shed their first flowers.
2. Rains occurring within 6 hours after spraying may reduce effectiveness. Should wash off occur within 6 hours, a single repeat application may be made.
3. Follow dosage recommendations carefully and spray field uniformly.
4. Compatibility studies with ROYAL MH-30SG and tobacco insecticides have not been completed. Separate application is recommended.
5. Do not use on tobacco plants grown for seed production.
6. Do not spray tobacco in the middle of hot, dry days when leaves are severely wilted. For best results spray during the early morning, all day on cloudy days, avoiding the late afternoon or night applications. (Use a nozzle delivering a fine mist spray.)
7. Closely hand-suckered tobacco or tobacco treated with ROYAL MH-30SG will not ripen as rapidly as poorly suckered tobacco. Also, under some conditions ROYAL MH-30SG treated tips have a tendency to turn yellow prematurely. This is not harmful and does not mean that your tobacco is ripe. Therefore, do not harvest tobacco treated with ROYAL MH-30SG by using time and color of the leaves as your only guide to ripeness. Use other characteristics indicative of ripeness such as:
  - a. Fading of color from midrib.
  - b. Change in apparent leaf texture.
8. If tobacco plants have been grown under drought conditions, absorption of ROYAL MH-30SG will be slow and results may not be as good as plants treated under normal conditions.
9. Too little ROYAL MH-30SG will result in less than the desired control, too much reduces quality of tips.
10. Use only one application of ROYAL MH-30SG per season.
11. Allow at least 7 days after application before harvest.
12. Even though ROYAL MH-30SG is non-volatile no drift should be permitted onto nearby crops.
13. Do not apply this product through any type of irrigation system.

## MIXING INSTRUCTIONS

1. Be sure sprayer is clean and not contaminated with any material.
2. Fill tank  $\frac{3}{4}$  full with clean water and start agitation.
3. Pour ROYAL MH-30SG slowly from bag into tank. Avoid dumping all contents at once.
4. Allow 2-3 minutes agitation for dissolving as tank filling is continued.

## TOBACCO

### DIRECTIONS FOR USE

To prevent sucker development on flue cured, burley, dark fire, Maryland and cigar (Pennsylvania, and Wisconsin) tobacco.

ROYAL MH-30SG is a potassium salt formulation of maleic hydrazide used to prevent sucker development in tobacco. Since ROYAL MH-30SG is absorbed and moves inside the tobacco plant, spraying the whole plant is not necessary. Best results are obtained by spraying the upper  $\frac{1}{2}$  to  $\frac{3}{4}$  of the plant. It is very important to apply equal amounts of ROYAL MH-30SG to each plant at the correct time to obtain satisfactory sucker control.

### TIMING

For flue cured, burley, dark fire and Maryland tobacco

1. If all plants flower at about the same time, proceed as follows: As soon as the plants are in EARLY FLOWER, top, pull all suckers from early maturing plants and spray ROYAL MH-30SG uniformly onto the entire field within 24 hours after topping. (See notes 1-12).

2. If there is a wide range in the time of flowering, the following procedure should be used:
  - a. Go through the field and top the few early flowering plants.
  - b. Wait until the remaining plants are in FULL FLOWER, top them, remove all suckers from the entire field, and apply ROYAL MH-30SG within 24 hours after topping. (See notes 1-12). Tip leaves to be harvested should be at least 6 to 8 inches long at time of spraying.
3. For best results ROYAL MH-30SG should be sprayed on dry or nearly dry leaves. Avoid spraying on leaves covered with dew.

### For Pennsylvania cigar tobacco

1. If your field flowers uniformly:
  - a. Top when about  $\frac{1}{4}$  to  $\frac{1}{2}$  of your plants show one or more open flowers on each head.
  - b. Wait three days after topping.
  - c. Spray ROYAL MH-30SG. For best results ROYAL MH-30SG should be sprayed on dry or nearly dry leaves. Avoid spraying on leaves covered with dew.
2. If your field doesn't flower uniformly:
  - a. When flowering in your field is particularly uneven you should top when about 50 to 60 percent of the plants show one or more open flowers per head.
  - b. Wait three days after topping and remove by hand any suckers more than two inches long.
  - c. Spray ROYAL MH-30SG over your entire field.

### For Wisconsin cigar tobacco

1. Top the field when most of the plants in your field are in full flower.
2. Remove, by hand, any suckers larger than one or two inches.
3. Spray within 24 hours after topping and removal of suckers.

## HOW TO SPRAY

ROYAL MH-30SG may be applied with either a conventional sprayer, or with a compressed air shoulder sprayer. Compressed air shoulder sprayer with nozzle spraying a wide cone: Spray one side of each tobacco row while walking at a constant rate. Use up and down motion of spray nozzle to moisten upper 6 to 8 leaves of each plant. Apply equal amounts of spray to each plant.

## RATE OF APPLICATION

For flue cured, burley, cigar

(Pennsylvania, Wisconsin) and Maryland Tobacco

Apply 4 to 5 pounds of ROYAL MH-30SG in 40 to 50 gallons of water per acre. Operate sprayer at 40 to 50 psi.

For dark tobacco

Apply 1 pound of ROYAL MH-30SG per 1000 plants. Use 40 to 50 gallons of water per acre and operate sprayer at 40 to 50 psi. Do not exceed 7.5 lbs. of ROYAL MH-30SG per acre.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a dry location.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**EQUIPMENT CLEANING:** ROYAL MH-30SG is not corrosive to spray equipment. Following applications, all spray equipment should be thoroughly rinsed with water.

**CONTAINER DISPOSAL:**

**Fiber carton with liners or bags:** Completely empty liner or bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner or bag in a sanitary landfill, or by incineration if permitted by State and local authorities. Dispose of carton in the same manner.

**BEST AVAILABLE COPY**

**IMPORTANT NOTICE**—Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

® ROYAL MH-30SG is a Registered Trademark of Uniroyal Chemical Company, Inc.

© Copyright 1993, Uniroyal Chemical Company, Inc.

**BEST AVAILABLE COPY**