

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS
WARNING**

Causes eye irritation. Do not get in eyes. Harmful if inhaled or absorbed through the skin. Avoid breathing spray mist. Avoid contact with skin or clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather favors drift from treated area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

EQUIPMENT CLEANING: Thoroughly rinse spray tank, hose and nozzles with water after application. Use RESCUE in metal or fiberglass tanks. RESCUE can be removed from metal or fiberglass tanks by washing thoroughly with ammonia water and rinsing several times with water containing a good detergent.

CONTAINER STORAGE AND HANDLING: RESCUE is normally a solution but it may crystallize in the cold (32 F). Gentle warming (60 F) will redissolve crystals. Mild agitation such as rolling, shaking or stirring before use is recommended.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

RESCUETM

POSTEMERGENT HERBICIDE

For Soybeans



COMPOSITION

Active Ingredient (percent by weight)	
Sodium N-1-naphthylphthalamate*	23.7%
Sodium salt of 4-(2,4-dichlorophenoxy) butyric acid**	6%
Inert Ingredients	75.7%
TOTAL	100.0%

*Equivalent to 2 lbs. of sodium N-1-naphthylphthalamate per gallon.

**Equivalent to 0.06 lbs 4-(2,4-DB) acid per gallon.

ACCEPTED

JAN 13 1983

U.S. Department of Agriculture
Insecticide, Fungicide, and Rodenticide Act,
and for the pesticide
registered under
EPA Reg. No. 400-166

KEEP OUT OF REACH OF CHILDREN.

WARNING

STATEMENTS OF PRACTICAL TREATMENT

If swallowed, drink two large glasses of water and induce vomiting. Contact a physician.
If on skin, wash affected area with plenty of soap and water.
If in eyes, flush eyes with plenty of water for at least 15 minutes. Contact a physician.
If inhaled, remove to fresh air; if abnormal symptoms develop, contact a physician.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Trademark of UNIROYAL, INC.
UNIROYAL CHEMICAL — Division of UNIROYAL, Inc.
Naugatuck, Connecticut 06770
EPA Reg. No. 400-
EPA Est. No.
NET CONTENTS:

LOT NO.

11-9-82

USE DIRECTIONS

GENERAL

RESCUE is a selective herbicide recommended for mid-to-late season postemergence weed control in soybeans. When applied according to label instructions, RESCUE will control or suppress the following broadleaf weeds, reducing their competition to the soybean crop: cocklebur, giant ragweed, sunflower (wild & volunteer), annual morningglory, and wild mustard.

Read and carefully observe all Application Notes prior to applying RESCUE.

APPLICATION TIMING

Determinate Soybeans - Apply when soybeans are at least 14 inches or taller.
Indeterminate Soybeans - Apply when soybeans are approximately 18 inches tall or after first bloom.

GROUND APPLICATION

Apply 2-3 quarts* of RESCUE in 10-25 gallons of water per acre (broadcast rate). A non-ionic spray adjuvant or crop oil concentrate at the rate of 2 quarts per 100 gallons of final spray solution should be used. The spray boom should be positioned 18-24" above the tops of the soybean plants or weeds and be equipped with hollow cone nozzles. Maintain a high pressure (40-50 p.s.i.) during application to insure better overall coverage. When making band applications, reduce rates and water volume according to width of band row spacing.

AERIAL APPLICATION

Apply 2-3 quarts* of RESCUE in a minimum of 5 gallons of water per acre. A non-ionic spray adjuvant or crop oil concentrate at the rate of 2 quarts per 100 gallons of final spray solution should be used.

Nozzle size and orientation should provide maximum droplet breakup without causing excessive drift.

*For best results, use the lower rate of RESCUE (2 quarts) on weeds 10 to 15 inches or less in height and the higher rate (3 quarts) if weeds are taller than 15 inches. Adequate control or suppression can only be expected if application is made before weeds reach 3 feet in height.