

**PRECAUTIONARY
STATEMENTS
HAZARDS TO HUMANS**

CAUTION

Harmful if swallowed. Avoid breathing of dust. Avoid contact with eyes or skin.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE — Store in a dry location. Do not store with propagative structures such as seed, bulbs, tubers, nursery stock, etc., or with food or feed products.

PESTICIDE DISPOSAL — Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.

CONTAINER DISPOSAL — Completely empty by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of container in a sanitary landfill or by incineration if allowed by State and local authorities.

BARRIER™

50W

DICHLOROBENIL WEED AND GRASS KILLER



COMPOSITION

Active ingredient: (% by weight)	
Dichlobenil*	50.0%
Inert Ingredients:	50.0%
TOTAL:	100.0%
*2,6-dichlorobenzonitrile, U.S. Patent 3,027,248	

ACCEPTED

1983

400-169

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See Side Panel For Additional Precautions

UNIROYAL CHEMICAL - Division of UNIROYAL, Inc.
NAUGATUCK, CONNECTICUT 06770
EPA Reg. No. 400-169
EPA Est. No.

2-24-83

GENERAL DIRECTIONS

ANNUAL WEED CONTROL

BARRIER 50W is recommended for control of annual grassy and broadleaved weeds and certain perennial weeds such as Annual Bluegrass (*Poa annua*), Bluegrass, Bull Thistle, Camphorweed, Carpetweed, Chickweed, Citron Melon, Cockweed, Crabgrass, Cudweed, Dandelion, Dogfennel, Evening Primrose, Fiddleneck, Florida Purslane (Pusley), Foxtail, Gisekia, Groundsel, Henbit, Horsetail, Jerusalem oak, Goosefoot, Knotweed, Lambsquarters, Maypops, Milkweed Vine (*Morrenia odorata*), Miners lettuce, Natalgrass, Old Witchgrass, Peppergrass, Pineapple Weed, Plantain, Purslane, Ragweed, Red Dead-nettle, Redroot Pigweed, Rosarypea, Russian Thistle, Shepherds-purse, Smartweed, Spanishneedles, Spurge, Tea-weed, Texas Panicum (Hurrahgrass), Wild Barley, Wild Mustard, Wild Radish, and Yellow Woodsorrel. Apply BARRIER 50W over-all or as a directed spray either in early spring before seeds of annual weeds germinate or after cultivation has removed the growing weeds. Apply in sufficient water for uniform coverage (at least 50 gallons per acre). If treatment is followed by application of 1/2 to 1 inch of water by overhead irrigation, the lower rates are recommended. Shallow incorporation immediately after application is recommended in southern areas. In other areas, shallow incorporation or sprinkler irrigation is suggested when applications are made during periods of high temperature. BARRIER 50W may be used for weed control on the plants listed below.

USE PRECAUTIONS

Do not use in seed beds, transplant or cutting beds or greenhouses. Do not apply until 6 months after rooting of cuttings in the field, or after planting rooted cuttings. Do not apply on fruit stock within 3 months prior to or following grafting or budding of root stocks or planting of new grafts. Do not apply more than the maximum recommended amount of BARRIER 50W per growing season. Do not use on "St. Lucie fine sand." "K-zell