

DEP ESTER

1064
PM 21
400-225

**SOYBEAN SEED
PROTECTANT**

highlighted
copy

UNIROYAL

ACCEPTED
SEP 18 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 400-225

FREE MEASURE INSIDE
(1/2 ounce capacity)

COMPOSITION
Active Ingredients: (% by weight)

Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	25%
Inert Ingredients:	75%
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN
DANGER
STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and see a physician.
IF INHALED: See a physician if an abnormal reaction occurs.
IF ON SKIN: Wash thoroughly with soap and water.
IF IN EYES: Immediately flush eyes with plenty of water and get medical treatment.
SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.



Uniroyal Chemical Company, Inc.
Middlebury, CT 06749
EPA REG. NO. 400 -225
EPA EST. NO.

5-11-86

NET WEIGHT:

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS**

WARNING

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

PHYSICAL or CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in dry location.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.


CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

- Slackwater Darter, Alabama (Lauderdale, Limestone and Madison)
Tennessee (Lawrence and Wayne)
- Pecos gambusia. New Mexico (Chaves and Eddy)
Texas (Reeves, Jeff Davis and Pecos)
- Freshwater Mussels Tennessee (Hancock Claiborne, Hawkins and Sullivan)
Virginia (Smyth, Scott, Lee, Washington and Grayson)
- Comanche Springs Pufffish, Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Mixers, loaders, and applicators, when mixing, loading, and applying, must wear mid-forearm to elbow-length natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan; boots or overshoes; one-piece overalls which have long sleeves and long pants; face shield or goggles and a hat or other appropriate head covering.

Fieldworkers and harvesters must wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan residues. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading his product.

Applicator's protection may also be obtained by use of an enclosed tractor cab with a properly filtered air supply.

Clothing worn while loading, mixing, and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

RECOMMENDATIONS

DE-PESTER SOYBEAN SEED PROTECTANT is especially formulated for treatment of soybean seed for the control of seed and soil-borne diseases, including damping-off and seed decay organisms, and seedling blights, and is prepared for use directly in the planter box, thereby eliminating additional seed handling to have seed treated. This material provides adequate coverage of seed and any excess flows with the seed into the furrow, giving additional protection against soil-borne diseases.

Do not use treated seed for food, feed or oil purposes.

Use DE-PESTER SOYBEAN SEED PROTECTANT at the rate of 4 oz. per bu. or 7 oz. per 100 lb. of seed as a planter box treatment. May be used on previously treated seed.

PLANTER BOX RATES ARE AS FOLLOWS:

Size of Planter Box	Number of measuring scoops per planter box
1/2 bushel	4 Heaping Scoops
1 bushel	8 Heaping Scoops

