

# DEPESTER®

MANEB, COPPER, SULFUR 4-5-15

**ACCEPTED**  
 JUL 11 1986  
 Registered under  
 EPA Reg. No. 400-243



### COMPOSITION

**ACTIVE INGREDIENTS: (% by weight)**

Maneb (Manganous ethylenebis-dithiocarbamate).....	4.0%
Copper expressed as metallic*...5.0%	
Sulfur.....	15.0%
<b>INERT INGREDIENTS.....</b>	<b>76.0%</b>
*Derived from Tribasic Copper Sulfate	<u>100.0%</u>

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**STATEMENT OF PRACTICAL TREATMENT**

"IF IN EYES, flush with plenty of water. Call a physician.  
 IF ON SKIN, wash with plenty of soap and water. Get medical attention. IF SWALLOWED, call a physician or...  
 Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or, if available, by administering syrup of ipecac. Do NOT induce vomiting or give anything by mouth to an unconscious person."

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT.

UNIROYAL CHEMICAL COMPANY, INC.  
 Middlebury, CT 06749  
 EPA REG. NO. 400-243  
 EPA EST. NO.

Lot No.

NET WEIGHT:

10/1

**DIRECTIONS FOR USE**

For application to either moist or dry foliage anytime dusting conditions are favorable. Application should begin when plants are small; increase dosage as plants grow larger. Dusters should be adjusted to give uniform and thorough coverage.

**RECOMMENDED USES**

**BEANS (Dry, Lima and String):** For control of downy mildew, powdery mildew, Anthracnose, and angular leaf spot, apply 15 to 35 lbs. per acre when mildew first appears and repeat at weekly intervals. Do not apply within 4 days of harvest.

**CARROTS:** For control of Cercospora leaf spot, Alternaria leaf blight, early blight, late blight and leaf spot, apply 15 to 45 lbs. per acre when diseases are reported in the area and repeat at 7 to 10 day intervals during the growing season or as required for adequate control.

**CELERY:** For control of early and late blight and leaf spot, apply 15 to 45 lbs. per acre when plants first come through ground. Repeat 7 to 10 day intervals, until plants are set in field. Continue field applications on weekly schedule. Remove excess residues by stripping, trimming, washing or other effective means before marketing.

**EGGPLANTS, PEPPERS, AND POTATOES:** For control of Anthracnose, and Cercosporae, early blight, late blight and leaf spot on Peppers: Phomopsis, early blight (alternaria) and anthracnose, early blight, late blight and leaf spot on Eggplants; early blight (Alternaria and late blight (Phytophthora) on potatoes, apply 25 to 45 lbs. per acre. Begin application when plants are transplanted to the field or before diseases are expected to appear, and repeat at 7 to 10 day intervals.


**ONIONS:** For control of downy mildew, botrytis leaf blight and purple blotch leaf blight and purple blotch, apply 25 to 50 lbs. per acre beginning when plants reach the 4 leaf stage or when disease conditions are reported in the area and repeat at 7 to 10 day intervals during season.

**SPINACH:** For control of downy mildew (blue mold), apply 25 to 45 lbs. per acre at first signs of mildew and repeat at 7 to 10 day intervals or as required. Do not apply within 7 days of harvest and remove excess residues by washing.

**TOMATOES:** For control of early and late blight, Anthracnose, Stemphylium (grey leaf spot) and Septoria leaf spots, apply 25 to 50 lbs. per acre when disease is first reported in area and repeat at 7 to 10 day intervals or oftener under severe late blight conditions. Wash and carefully wipe all fresh fruit before marketing.

Do not use on Cucurbits or other plants susceptible to sulfur injury. Store in a cool, dry place. Keep container closed when not in use. Harmful if swallowed. Copper 5% dust retains toxicity for a long period of time when exposed to weathering action. This product is toxic to fungus diseases, but will not injure fruit or vegetables when used at recommended dosages.

Consult your local agricultural authorities for further details as to time, frequency and rate of applications.



**IMPORTANT NOTICE** — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

2.5/1

### REENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written or oral warnings must include the following information:

**"WARNING"** "Area to be treated with ROYAL MH-30SG on (date of application). Do not enter without appropriate protective clothing until sprays have dried. If accidentally exposed, consult the Statement of Practical Treatment portion of the pesticide label for first aid recommendations."

### "ENDANGERED SPECIES RESTRICTIONS"

It is a violation of Federal laws to use any pesticide in a manner that results in the death of an endangered species or adverse modification of their habitat.

The use of this product may pose a hazard to certain Federally designated endangered species known to occur in specific areas within the following counties:

STATE	COUNTY
Species (Bulletin No.)	
CALIFORNIA Solano Grass (EPA/ES-85-13)	Solano
TENNESSEE Slackwater Darter (EPA/ES-85-04)	Lawrence Wayne Hancock
Freshwater Mussels (EPA/ES-85-07)	Claiborne Hawkins Sullivan
ALABAMA Slackwater Darter (EPA/ES-85-05)	Lauderdale Limestone Madison
VIRGINIA Freshwater Mussels (EPA/ES-85-06)	Grayson Smyth Scott Lee Washington

Before using this product in the above counties you must obtain the EPA Bulletin specific to your area. This Bulletin identifies areas within these counties where the use of this pesticide is prohibited, unless specified otherwise. The EPA Bulletin is available from either your County Agricultural Extension Agent, the Endangered Species Specialist in your State Wildlife Agency Headquarters, or the appropriate Regional Office of the U.S. Fish and Wildlife Service. THIS BULLETIN MUST BE REVIEWED PRIOR TO PESTICIDE USE.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS**

**CAUTION**

May cause irritation of nose and throat. Do not breathe dust. May cause skin sensitization in certain individuals.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

STORAGE-Store in a dry location.

PESTICIDE DISPOSAL-Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility.

CONTAINER DISPOSAL - Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.



1/0/4